

第3部

身体状況調査の結果

第8表 身長・体重の平均値及び標準偏差（1歳以上、性・年齢階級別）

年齢(歳)	男性						女性					
	身長(cm)			体重(kg)			身長(cm)			体重(kg)		
	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差
総数	206	162.0	5.2	206	61.5	9.1	319	150.9	5.6	318	51.2	8.6
1	-	-	-	-	-	-	1	72.0	-	1	8.7	-
2	2	84.5	3.8	2	12.9	2.0	-	-	-	-	-	-
3	1	98.2	-	1	15.7	-	-	-	-	-	-	-
4	2	102.2	0.3	2	15.7	0.6	3	99.8	0.9	3	14.8	2.6
5	-	-	-	-	-	-	-	-	-	-	-	-
6	-	-	-	-	-	-	1	119.5	-	1	21.5	-
7	4	120.4	2.4	4	22.5	2.4	-	-	-	-	-	-
8	1	117.0	-	1	19.9	-	4	127.6	1.2	4	26.9	3.5
9	2	130.1	6.4	2	25.3	3.4	2	131.1	0.3	2	24.2	0.6
10	2	131.6	2.0	2	28.9	1.3	1	135.7	-	1	26.8	-
11	2	142.5	2.3	2	33.9	1.8	1	145.0	-	1	37.0	-
12	1	154.7	-	1	38.5	-	5	153.7	3.4	5	42.1	4.6
13	-	-	-	-	-	-	1	152.0	-	1	40.0	-
14	1	158.5	-	1	44.1	-	1	160.0	-	1	52.0	-
15	4	167.9	2.5	4	57.0	11.0	1	154.0	-	1	47.0	-
16	-	-	-	-	-	-	2	153.3	4.1	2	44.6	0.5
17	1	170.0	-	1	54.0	-	-	-	-	-	-	-
18	2	170.3	1.8	2	70.0	12.6	-	-	-	-	-	-
19	-	-	-	-	-	-	2	157.0	2.9	2	48.1	0.6
20	-	-	-	-	-	-	-	-	-	-	-	-
21	-	-	-	-	-	-	1	160.7	-	1	46.8	-
22	-	-	-	-	-	-	2	163.7	4.1	2	50.3	6.2
23	1	182.2	-	1	84.3	-	2	148.3	1.1	2	33.7	1.6
24	-	-	-	-	-	-	-	-	-	-	-	-
25	-	-	-	-	-	-	3	160.4	6.3	3	58.4	18.6
26-29	5	174.5	5.4	5	64.1	7.5	5	156.7	3.9	5	59.2	9.3
30-39	12	169.8	6.6	12	67.6	9.2	13	159.4	5.9	13	55.5	6.8
40-49	17	170.7	5.1	17	67.6	9.9	27	158.3	5.0	26	55.2	6.8
50-59	22	167.8	4.9	22	63.4	9.0	43	157.4	6.4	43	55.2	9.0
60-69	48	167.2	5.2	48	67.5	10.2	75	152.6	6.7	75	54.3	9.6
70歳以上	76	161.6	6.2	76	63.3	7.8	123	147.5	6.4	123	50.2	8.8
(再掲)												
20歳以上	181	165.7	6.8	181	65.3	9.1	294	152.2	7.8	293	52.8	9.4
20-29歳	6	175.8	5.8	6	67.5	10.7	13	157.6	6.5	13	52.8	14.6
60-64歳	20	169.2	4.6	20	67.2	12.4	31	153.4	6.5	31	54.7	9.7
65-69歳	28	165.8	5.1	28	67.8	8.5	44	152.1	6.9	44	54.0	9.6
70-74歳	24	163.7	5.4	24	62.1	6.1	42	150.8	5.8	42	50.5	8.0
75-79歳	25	161.5	5.6	25	65.0	8.1	42	147.4	6.1	42	52.2	9.3
80歳以上	27	160.0	7.0	27	62.9	8.8	39	143.9	5.4	39	47.7	8.8

注) 体重は妊婦除外

第9表 BMIの平均値及び標準偏差（15歳以上、性・年齢階級別）

年齢(歳)	男性			女性		
	人数	平均値	標準偏差	人数	平均値	標準偏差
総数	188	23.7	3.6	298	22.7	4.1
15-19歳	7	21.1	4.5	5	19.4	1.0
20-29歳	6	21.8	3.0	13	21.1	5.8
30-39歳	12	23.6	4.2	13	21.8	1.6
40-49歳	17	23.2	3.4	26	22.1	3.6
50-59歳	22	22.6	3.4	43	22.3	3.5
60-69歳	48	24.1	3.4	75	23.3	4.3
70歳以上	76	24.3	2.8	123	23.1	4.2

注) 妊婦除外

第10表 BMIの分布（15歳以上、男女別）

年齢(歳)	総数		男性		女性	
	人数	%	人数	%	人数	%
総数	486	100.0	188	100.0	298	100.0
15未満	2	0.4	-	-	2	0.7
15-16未満	5	1.0	1	0.5	4	1.3
16-17未満	5	1.0	-	-	5	1.7
17-18未満	14	2.9	5	2.7	9	3.0
18-19未満	26	5.3	8	4.3	18	6.0
19-20未満	44	9.1	9	4.8	35	11.7
20-21未満	49	10.1	15	8.0	34	11.4
21-22未満	40	8.2	18	9.6	22	7.4
22-23未満	72	14.8	22	11.7	50	16.8
23-24未満	47	9.7	24	12.8	23	7.7
24-25未満	48	9.9	20	10.6	28	9.4
25-26未満	40	8.2	27	14.4	13	4.4
26-27未満	37	7.6	18	9.6	19	6.4
27-28未満	17	3.5	6	3.2	11	3.7
28-29未満	7	1.4	3	1.6	4	1.3
29-30未満	13	2.7	5	2.7	8	2.7
30-31未満	4	0.8	1	0.5	3	1.0
31-32未満	5	1.0	2	1.1	3	1.0
32-33未満	3	0.6	1	0.5	2	0.7
33-34未満	3	0.6	1	0.5	2	0.7
34-35未満	1	0.2	1	0.5	-	-
35以上	4	0.8	1	0.5	3	1.0

注) 妊婦除外

第11表 BMIの状況(15歳以上、性・年齢階級別)

		総数		肥満区分					
				やせ		普通		肥満	
		人数	%	人数	%	人数	%	人数	%
男 性	総数	188	100.0	8	4.3	114	60.6	66	35.1
	15-19歳	7	100.0	2	28.6	3	42.9	2	28.6
	20-29歳	6	100.0	-	-	4	66.7	2	33.3
	30-39歳	12	100.0	1	8.3	9	75.0	2	16.7
	40-49歳	17	100.0	-	-	13	76.5	4	23.5
	50-59歳	22	100.0	3	13.6	14	63.6	5	22.7
	60-69歳	48	100.0	1	2.1	29	60.4	18	37.5
	70歳以上	76	100.0	1	1.3	42	55.3	33	43.4
	(再掲)								
	20歳以上	181	100.0	6	3.3	111	61.3	64	35.4
	20-39歳	18	100.0	1	5.6	13	72.2	4	22.2
	20-69歳	105	100.0	5	4.8	69	65.7	31	29.5
	40-69歳	87	100.0	4	4.6	56	64.4	27	31.0
	65-69歳	28	100.0	-	-	15	53.6	13	46.4
	70-74歳	24	100.0	-	-	17	70.8	7	29.2
75-79歳	25	100.0	-	-	11	44.0	14	56.0	
80-84歳	19	100.0	-	-	10	52.6	9	47.4	
85歳以上	8	100.0	1	12.5	4	50.0	3	37.5	
女 性	総数	298	100.0	26	8.7	204	68.5	68	22.8
	15-19歳	5	100.0	1	20.0	4	80.0	-	-
	20-29歳	13	100.0	5	38.5	5	38.5	3	23.1
	30-39歳	13	100.0	1	7.7	12	92.3	-	-
	40-49歳	26	100.0	-	-	23	88.5	3	11.5
	50-59歳	43	100.0	4	9.3	32	74.4	7	16.3
	60-69歳	75	100.0	5	6.7	45	60.0	25	33.3
	70歳以上	123	100.0	10	8.1	83	67.5	30	24.4
	(再掲)								
	20歳以上	293	100.0	25	8.5	200	68.3	68	23.2
	20-39歳	26	100.0	6	23.1	17	65.4	3	11.5
	20-69歳	172	100.0	17	9.9	117	68.0	38	22.1
	40-69歳	144	100.0	9	6.3	100	69.4	35	24.3
	65-69歳	44	100.0	2	4.5	26	59.1	16	36.4
	70-74歳	42	100.0	6	14.3	31	73.8	5	11.9
75-79歳	42	100.0	1	2.4	26	61.9	15	35.7	
80-84歳	20	100.0	3	15.0	12	60.0	5	25.0	
85歳以上	19	100.0	-	-	14	73.7	5	26.3	

注) 妊婦除外 BMIによる肥満の判定は p.5 参照

第12表 低栄養傾向（BMI20以下）の高齢者の割合（65歳以上、男女別）

		総数		BMI 20未満		BMI 20以上	
		人数	%	人数	%	人数	%
総数	65歳以上	271	100.0	41	15.1	230	84.9
	（再掲）75歳以上	133	100.0	18	13.5	115	86.5
男性	65歳以上	104	100.0	6	5.8	98	94.2
	（再掲）75歳以上	52	100.0	4	7.7	48	92.3
女性	65歳以上	167	100.0	35	21.0	132	79.0
	（再掲）75歳以上	81	100.0	14	17.3	67	82.7

第13表 日比式による肥満区分（6-14歳、性・年齢階級別）

		総数		6-8歳		9-11歳		12-14歳	
		人数	%	人数	%	人数	%	人数	%
総数	総数	29	100.0	10	100.0	10	100.0	9	100.0
	-30%未満	1	3.4	-	-	-	-	1	11.1
	-30%以上-20%未満	-	-	-	-	-	-	-	-
	-20%以上-10%未満	8	27.6	2	20.0	3	30.0	3	33.3
	-10%以上0%未満	11	37.9	4	40.0	6	60.0	1	11.1
	0%以上10%未満	5	17.2	3	30.0	-	-	4	44.4
	10%以上20%未満	3	10.3	-	-	1	10.0	-	-
	20%以上30%未満	1	3.4	1	10.0	-	-	-	-
	30%以上40%未満	-	-	-	-	-	-	-	-
	40%以上	-	-	-	-	-	-	-	-
男性	総数	13	100.0	5	100.0	6	100.0	2	100.0
	-30%未満	-	-	-	-	-	-	-	-
	-30%以上-20%未満	-	-	-	-	-	-	-	-
	-20%以上-10%未満	3	23.1	1	20.0	-	-	2	100.0
	-10%以上0%未満	7	53.8	2	40.0	5	83.3	-	-
	0%以上10%未満	2	15.4	2	40.0	-	-	-	-
	10%以上20%未満	1	7.7	-	-	1	16.7	-	-
	20%以上30%未満	-	-	-	-	-	-	-	-
	30%以上40%未満	-	-	-	-	-	-	-	-
	40%以上	-	-	-	-	-	-	-	-
女性	総数	16	100.0	5	100.0	4	100.0	7	100.0
	-30%未満	1	6.3	-	-	-	-	1	14.3
	-30%以上-20%未満	-	-	-	-	-	-	-	-
	-20%以上-10%未満	5	31.3	1	20.0	3	75.0	1	14.3
	-10%以上0%未満	4	25.0	2	40.0	1	25.0	1	14.3
	0%以上10%未満	3	18.8	1	20.0	-	-	4	57.1
	10%以上20%未満	2	12.5	-	-	-	-	-	-
	20%以上30%未満	1	6.3	1	20.0	-	-	-	-
	30%以上40%未満	-	-	-	-	-	-	-	-
	40%以上	-	-	-	-	-	-	-	-

注) 日比式による区分は p.5 参照

第14表 学校保健統計調査方式による肥満区分（6-14歳、性・年齢階級別）

		総数		6-8歳		9-11歳		12-14歳	
		人数	%	人数	%	人数	%	人数	%
総 数	総数	29	100.0	10	100.0	10	100.0	9	100.0
	-30%以下	1	3.4	-	-	-	-	1	11.1
	-30%超-20%以下	-	-	-	-	-	-	-	-
	-20%超-10%以下	7	24.1	2	20.0	4	40.0	1	11.1
	-10%超0%以下	12	41.4	4	40.0	5	50.0	3	33.3
	0%超10%以下	4	13.8	3	30.0	-	-	1	11.1
	10%超20%以下	5	17.2	1	10.0	1	10.0	3	33.3
	20%超30%以下	-	-	-	-	-	-	-	-
	30%超40%以下	-	-	-	-	-	-	-	-
40%超	-	-	-	-	-	-	-	-	
男 性	総数	13	100.0	5	100.0	6	100.0	2	100.0
	-30%以下	-	-	-	-	-	-	-	-
	-30%超-20%以下	-	-	-	-	-	-	-	-
	-20%超-10%以下	3	23.1	1	20.0	1	16.7	1	50.0
	-10%超0%以下	7	53.8	2	40.0	4	66.7	1	50.0
	0%超10%以下	2	15.4	2	40.0	-	-	-	-
	10%超20%以下	1	7.7	-	-	1	16.7	-	-
	20%超30%以下	-	-	-	-	-	-	-	-
	30%超40%以下	-	-	-	-	-	-	-	-
40%超	-	-	-	-	-	-	-	-	
女 性	総数	16	100.0	5	100.0	4	100.0	7	100.0
	-30%以下	1	6.3	-	-	-	-	1	14.3
	-30%超-20%以下	-	-	-	-	-	-	-	-
	-20%超-10%以下	4	25.0	1	20.0	3	75.0	-	-
	-10%超0%以下	5	31.3	2	40.0	1	25.0	2	28.6
	0%超10%以下	2	12.5	1	20.0	-	-	1	14.3
	10%超20%以下	4	25.0	1	20.0	-	-	3	42.9
	20%超30%以下	-	-	-	-	-	-	-	-
	30%超40%以下	-	-	-	-	-	-	-	-
40%超	-	-	-	-	-	-	-	-	

注) 学校保健統計調査方式による区分は p.6 参照

第15表 腹囲の分布（20歳以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	40-74歳	
															人数	%
総数	441	100.0	19	100.0	23	100.0	37	100.0	64	100.0	117	100.0	181	100.0	280	100.0
55.0未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55.0～59.9	1	0.2	1	5.3	-	-	-	-	-	-	-	-	-	-	-	
60.0～64.9	12	2.7	3	15.8	-	-	1	2.7	1	1.6	2	1.7	5	2.8	8	2.9
65.0～69.9	22	5.0	4	21.1	1	4.3	-	-	5	7.8	4	3.4	8	4.4	11	3.9
70.0～74.9	46	10.4	2	10.5	4	17.4	9	24.3	8	12.5	11	9.4	12	6.6	33	11.8
75.0～79.9	53	12.0	4	21.1	5	21.7	4	10.8	12	18.8	17	14.5	11	6.1	37	13.2
80.0～84.9	96	21.8	-	-	9	39.1	11	29.7	11	17.2	16	13.7	49	27.1	55	19.6
85.0～89.9	89	20.2	2	10.5	2	8.7	7	18.9	11	17.2	24	20.5	43	23.8	58	20.7
90.0～94.9	73	16.6	1	5.3	1	4.3	3	8.1	11	17.2	26	22.2	31	17.1	52	18.6
95.0～99.9	25	5.7	2	10.5	-	-	1	2.7	3	4.7	10	8.5	9	5.0	15	5.4
100.0～104.9	15	3.4	-	-	1	4.3	1	2.7	2	3.1	3	2.6	8	4.4	7	2.5
105.0～109.9	7	1.6	-	-	-	-	-	-	-	-	2	1.7	5	2.8	2	0.7
110.0～114.9	1	0.2	-	-	-	-	-	-	-	-	1	0.9	-	-	1	0.4
115.0～119.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0～124.9	1	0.2	-	-	-	-	-	-	-	-	1	0.9	-	-	1	0.4
125.0以上	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
総数	160	100.0	6	100.0	10	100.0	13	100.0	22	100.0	44	100.0	65	100.0	99	100.0
55.0未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55.0～59.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
60.0～64.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65.0～69.9	1	0.6	-	-	-	-	-	-	1	4.5	-	-	-	-	1	1.0
70.0～74.9	10	6.3	1	16.7	1	10.0	2	15.4	4	18.2	2	4.5	-	-	8	8.1
75.0～79.9	20	12.5	3	50.0	2	20.0	1	7.7	5	22.7	6	13.6	3	4.6	14	14.1
80.0～84.9	34	21.3	-	-	4	40.0	3	23.1	5	22.7	6	13.6	16	24.6	21	21.2
85.0～89.9	41	25.6	1	16.7	1	10.0	3	23.1	3	13.6	10	22.7	23	35.4	24	24.2
90.0～94.9	41	25.6	1	16.7	1	10.0	3	23.1	3	13.6	14	31.8	19	29.2	23	23.2
95.0～99.9	4	2.5	-	-	-	-	-	-	-	-	2	4.5	2	3.1	2	2.0
100.0～104.9	5	3.1	-	-	1	10.0	1	7.7	1	4.5	2	4.5	-	-	4	4.0
105.0～109.9	3	1.9	-	-	-	-	-	-	-	-	1	2.3	2	3.1	1	1.0
110.0～114.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
115.0～119.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0～124.9	1	0.6	-	-	-	-	-	-	-	-	1	2.3	-	-	1	1.0
125.0以上	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
総数	281	100.0	13	100.0	13	100.0	24	100.0	42	100.0	73	100.0	116	100.0	181	100.0
55.0未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
55.0～59.9	1	0.4	1	7.7	-	-	-	-	-	-	-	-	-	-	-	-
60.0～64.9	12	4.3	3	23.1	-	-	1	4.2	1	2.4	2	2.7	5	4.3	8	4.4
65.0～69.9	21	7.5	4	30.8	1	7.7	-	-	4	9.5	4	5.5	8	6.9	10	5.5
70.0～74.9	36	12.8	1	7.7	3	23.1	7	29.2	4	9.5	9	12.3	12	10.3	25	13.8
75.0～79.9	33	11.7	1	7.7	3	23.1	3	12.5	7	16.7	11	15.1	8	6.9	23	12.7
80.0～84.9	62	22.1	-	-	5	38.5	8	33.3	6	14.3	10	13.7	33	28.4	34	18.8
85.0～89.9	48	17.1	1	7.7	1	7.7	4	16.7	8	19.0	14	19.2	20	17.2	34	18.8
90.0～94.9	32	11.4	-	-	-	-	-	-	8	19.0	12	16.4	12	10.3	29	16.0
95.0～99.9	21	7.5	2	15.4	-	-	1	4.2	3	7.1	8	11.0	7	6.0	13	7.2
100.0～104.9	10	3.6	-	-	-	-	-	-	1	2.4	1	1.4	8	6.9	3	1.7
105.0～109.9	4	1.4	-	-	-	-	-	-	-	-	1	1.4	3	2.6	1	0.6
110.0～114.9	1	0.4	-	-	-	-	-	-	-	-	1	1.4	-	-	1	0.6
115.0～119.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
120.0～124.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
125.0以上	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

注) 妊婦除外

第16表 腹囲の区分（20歳以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	441	100.0	19	100.0	23	100.0	37	100.0	64	100.0	117	100.0	181	100.0	280	100.0
85cm未満(男)/90cm未満(女)	278	63.0	15	78.9	20	87.0	29	78.4	45	70.3	64	54.7	105	58.0	178	63.6
85cm以上(男)/90cm以上(女)	163	37.0	4	21.1	3	13.0	8	21.6	19	29.7	53	45.3	76	42.0	102	36.4
総数	160	100.0	6	100.0	10	100.0	13	100.0	22	100.0	44	100.0	65	100.0	99	100.0
男性	65	40.6	4	66.7	7	70.0	6	46.2	15	68.2	14	31.8	19	29.2	44	44.4
85cm未満	95	59.4	2	33.3	3	30.0	7	53.8	7	31.8	30	68.2	46	70.8	55	55.6
85cm以上	281	100.0	13	100.0	13	100.0	24	100.0	42	100.0	73	100.0	116	100.0	181	100.0
女性	213	75.8	11	84.6	13	100.0	23	95.8	30	71.4	50	68.5	86	74.1	134	74.0
85cm未満	68	24.2	2	15.4	-	-	1	4.2	12	28.6	23	31.5	30	25.9	47	26.0
85cm以上																

注) 妊婦除外

第17表 BMI・腹囲の区分（20歳以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲) 40-74歳	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	441	100.0	19	100.0	23	100.0	37	100.0	64	100.0	117	100.0	181	100.0	280	100.0
BMI25未満、85cm未満(男)/90cm未満(女)	262	59.4	14	73.7	20	87.0	27	73.0	44	68.8	57	48.7	100	55.2	168	60.0
BMI25未満、85cm以上(男)/90cm以上(女)	58	13.2	-	-	1	4.3	3	8.1	9	14.1	20	17.1	25	13.8	42	15.0
BMI25以上、85cm未満(男)/90cm未満(女)	16	3.6	1	5.3	-	-	2	5.4	1	1.6	7	6.0	5	2.8	10	3.6
BMI25以上、85cm以上(男)/90cm以上(女)	105	23.8	4	21.1	2	8.7	5	13.5	10	15.6	33	28.2	51	28.2	60	21.4
総数	160	100.0	6	100.0	10	100.0	13	100.0	22	100.0	44	100.0	65	100.0	99	100.0
男性	62	38.8	4	66.7	7	70.0	6	46.2	14	63.6	14	31.8	17	26.2	43	43.4
BMI25未満、腹囲85cm未満	38	23.8	-	-	1	10.0	3	23.1	3	13.6	13	29.5	18	27.7	23	23.2
BMI25未満、腹囲85cm以上	3	1.9	-	-	-	-	-	-	1	4.5	-	-	2	3.1	1	1.0
BMI25以上、腹囲85cm未満	57	35.6	2	33.3	2	20.0	4	30.8	4	18.2	17	38.6	28	43.1	32	32.3
BMI25以上、腹囲85cm以上	281	100.0	13	100.0	13	100.0	24	100.0	42	100.0	73	100.0	116	100.0	181	100.0
女性	200	71.2	10	76.9	13	100.0	21	87.5	30	71.4	43	58.9	83	71.6	125	69.1
BMI25未満、腹囲90cm未満	20	7.1	-	-	-	-	-	-	6	14.3	7	9.6	7	6.0	19	10.5
BMI25未満、腹囲90cm以上	13	4.6	1	7.7	-	-	2	8.3	-	-	7	9.6	3	2.6	9	5.0
BMI25以上、腹囲90cm未満	48	17.1	2	15.4	-	-	1	4.2	6	14.3	16	21.9	23	19.8	28	15.5
BMI25以上、腹囲90cm以上																

注) 妊婦除外

第18表 糖尿病の指摘の状況（20歳以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)					
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	40-74歳		65-74歳		75歳以上			
													人数	%	人数	%	人数	%	人数	%
総数	435	100.0	19	100.0	22	100.0	37	100.0	64	100.0	116	100.0	177	100.0	279	100.0	130	100.0	115	100.0
あり	69	15.9	-	-	-	-	-	-	4	6.3	27	23.3	38	21.5	44	15.8	28	21.5	25	21.7
なし	366	84.1	19	100.0	22	100.0	37	100.0	60	93.8	89	76.7	139	78.5	235	84.2	102	78.5	90	78.3
総数	156	100.0	6	100.0	10	100.0	13	100.0	22	100.0	44	100.0	61	100.0	99	100.0	47	100.0	41	100.0
あり	35	22.4	-	-	-	-	-	-	3	13.6	16	36.4	16	26.2	24	24.2	15	31.9	11	26.8
なし	121	77.6	6	100.0	10	100.0	13	100.0	19	86.4	28	63.6	45	73.8	75	75.8	32	68.1	30	73.2
総数	279	100.0	13	100.0	12	100.0	24	100.0	42	100.0	72	100.0	116	100.0	180	100.0	83	100.0	74	100.0
あり	34	12.2	-	-	-	-	-	-	1	2.4	11	15.3	22	19.0	20	11.1	13	15.7	14	18.9
なし	245	87.8	13	100.0	12	100.0	24	100.0	41	97.6	61	84.7	94	81.0	160	88.9	70	84.3	60	81.1

身体状況調査票(7)これまでに医療機関や健診で糖尿病といわれたことの有無(「境界型である」、「糖尿病の気がある」、「糖尿病になりかけている」、「血糖値が高い」などのように言われたことも含む)

第19表 糖尿病を指摘されたことのある者における治療の状況（20歳以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)					
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	40-74歳		65-74歳		75歳以上			
													人数	%	人数	%	人数	%	人数	%
総数	69	100.0	-	-	-	-	-	-	4	100.0	27	100.0	38	100.0	44	100.0	28	100.0	25	100.0
あり	43	62.3	-	-	-	-	-	-	1	25.0	16	59.3	26	68.4	26	59.1	18	64.3	17	68.0
なし	26	37.7	-	-	-	-	-	-	3	75.0	11	40.7	12	31.6	18	40.9	10	35.7	8	32.0
総数	35	100.0	-	-	-	-	-	-	3	100.0	16	100.0	16	100.0	24	100.0	15	100.0	11	100.0
あり	24	68.6	-	-	-	-	-	-	1	33.3	10	62.5	13	81.3	15	62.5	11	73.3	9	81.8
なし	11	31.4	-	-	-	-	-	-	2	66.7	6	37.5	3	18.8	9	37.5	4	26.7	2	18.2
総数	34	100.0	-	-	-	-	-	-	1	100.0	11	100.0	22	100.0	20	100.0	13	100.0	14	100.0
あり	19	55.9	-	-	-	-	-	-	-	-	6	54.5	13	59.1	11	55.0	7	53.8	8	57.1
なし	15	44.1	-	-	-	-	-	-	1	100.0	5	45.5	9	40.9	9	45.0	6	46.2	6	42.9

身体状況調査票(7-1) 現在、糖尿病の治療の有無(通院による定期的な検査や生活習慣の改善指導を含む)

注)(7)で1(有)と回答した者

第20表の1 血圧を下げる薬の服用状況
(20歳以上、性・年齢階級別)

		総数		有		無	
		人数	%	人数	%	人数	%
総数	総数	439	100.0	138	31.4	301	68.6
	20-29歳	19	100.0	-	-	19	100.0
	30-39歳	22	100.0	-	-	22	100.0
	40-49歳	37	100.0	-	-	37	100.0
	50-59歳	64	100.0	12	18.8	52	81.3
	60-69歳	116	100.0	34	29.3	82	70.7
	70歳以上	181	100.0	92	50.8	89	49.2
男性	総数	159	100.0	59	37.1	100	62.9
	20-29歳	6	100.0	-	-	6	100.0
	30-39歳	10	100.0	-	-	10	100.0
	40-49歳	13	100.0	-	-	13	100.0
	50-59歳	22	100.0	7	31.8	15	68.2
	60-69歳	44	100.0	16	36.4	28	63.6
	70歳以上	64	100.0	36	56.3	28	43.8
女性	総数	280	100.0	79	28.2	201	71.8
	20-29歳	13	100.0	-	-	13	100.0
	30-39歳	12	100.0	-	-	12	100.0
	40-49歳	24	100.0	-	-	24	100.0
	50-59歳	42	100.0	5	11.9	37	88.1
	60-69歳	72	100.0	18	25.0	54	75.0
	70歳以上	117	100.0	56	47.9	61	52.1

注) 妊婦除外

第20表の2 脈の乱れを治す薬の服用状況
(20歳以上、性・年齢階級別)

		総数		有		無	
		人数	%	人数	%	人数	%
総数	総数	438	100.0	17	3.9	421	96.1
	20-29歳	19	100.0	-	-	19	100.0
	30-39歳	22	100.0	-	-	22	100.0
	40-49歳	37	100.0	-	-	37	100.0
	50-59歳	64	100.0	3	4.7	61	95.3
	60-69歳	116	100.0	5	4.3	111	95.7
	70歳以上	180	100.0	9	5.0	171	95.0
男性	総数	159	100.0	9	5.7	150	94.3
	20-29歳	6	100.0	-	-	6	100.0
	30-39歳	10	100.0	-	-	10	100.0
	40-49歳	13	100.0	-	-	13	100.0
	50-59歳	22	100.0	1	4.5	21	95.5
	60-69歳	44	100.0	2	4.5	42	95.5
	70歳以上	64	100.0	6	9.4	58	90.6
女性	総数	279	100.0	8	2.9	271	97.1
	20-29歳	13	100.0	-	-	13	100.0
	30-39歳	12	100.0	-	-	12	100.0
	40-49歳	24	100.0	-	-	24	100.0
	50-59歳	42	100.0	2	4.8	40	95.2
	60-69歳	72	100.0	3	4.2	69	95.8
	70歳以上	116	100.0	3	2.6	113	97.4

注) 妊婦除外

第20表の3 インスリン注射または血糖を下げる薬の服用状況 (20歳以上、性・年齢階級別)

		総数		有		無	
		人数	%	人数	%	人数	%
総数	総数	438	100.0	27	6.2	411	93.8
	20-29歳	19	100.0	-	-	19	100.0
	30-39歳	22	100.0	-	-	22	100.0
	40-49歳	37	100.0	-	-	37	100.0
	50-59歳	64	100.0	-	-	64	100.0
	60-69歳	116	100.0	9	7.8	107	92.2
	70歳以上	180	100.0	18	10.0	162	90.0
男性	総数	158	100.0	12	7.6	146	92.4
	20-29歳	6	100.0	-	-	6	100.0
	30-39歳	10	100.0	-	-	10	100.0
	40-49歳	13	100.0	-	-	13	100.0
	50-59歳	22	100.0	-	-	22	100.0
	60-69歳	44	100.0	5	11.4	39	88.6
	70歳以上	63	100.0	7	11.1	56	88.9
女性	総数	280	100.0	15	5.4	265	94.6
	20-29歳	13	100.0	-	-	13	100.0
	30-39歳	12	100.0	-	-	12	100.0
	40-49歳	24	100.0	-	-	24	100.0
	50-59歳	42	100.0	-	-	42	100.0
	60-69歳	72	100.0	4	5.6	68	94.4
	70歳以上	117	100.0	11	9.4	106	90.6

注) 妊婦除外

第20表の4 コレステロールを下げる薬の服用状況 (20歳以上、性・年齢階級別)

		総数		有		無	
		人数	%	人数	%	人数	%
総数	総数	436	100.0	89	20.4	347	79.6
	20-29歳	19	100.0	-	-	19	100.0
	30-39歳	22	100.0	-	-	22	100.0
	40-49歳	37	100.0	-	-	37	100.0
	50-59歳	64	100.0	10	15.6	54	84.4
	60-69歳	115	100.0	31	27.0	84	73.0
	70歳以上	179	100.0	48	26.8	131	73.2
男性	総数	157	100.0	23	14.6	134	85.4
	20-29歳	6	100.0	-	-	6	100.0
	30-39歳	10	100.0	-	-	10	100.0
	40-49歳	13	100.0	-	-	13	100.0
	50-59歳	22	100.0	3	13.6	19	86.4
	60-69歳	44	100.0	7	15.9	37	84.1
	70歳以上	62	100.0	13	21.0	49	79.0
女性	総数	279	100.0	66	23.7	213	76.3
	20-29歳	13	100.0	-	-	13	100.0
	30-39歳	12	100.0	-	-	12	100.0
	40-49歳	24	100.0	-	-	24	100.0
	50-59歳	42	100.0	7	16.7	35	83.3
	60-69歳	71	100.0	24	33.8	47	66.2
	70歳以上	117	100.0	35	29.9	82	70.1

注) 妊婦除外

第 20 表の 5 中性脂肪（トリグリセライド）を
下げる薬の服用状況（20 歳以上、性・年齢階級別）

		総数		有		無	
		人数	%	人数	%	人数	%
総 数	総数	436	100.0	29	6.7	407	93.3
	20-29歳	19	100.0	-	-	19	100.0
	30-39歳	22	100.0	-	-	22	100.0
	40-49歳	37	100.0	-	-	37	100.0
	50-59歳	64	100.0	4	6.3	60	93.8
	60-69歳	115	100.0	10	8.7	105	91.3
	70歳以上	179	100.0	15	8.4	164	91.6
男 性	総数	157	100.0	12	7.6	145	92.4
	20-29歳	6	100.0	-	-	6	100.0
	30-39歳	10	100.0	-	-	10	100.0
	40-49歳	13	100.0	-	-	13	100.0
	50-59歳	22	100.0	3	13.6	19	86.4
	60-69歳	44	100.0	5	11.4	39	88.6
	70歳以上	62	100.0	4	6.5	58	93.5
女 性	総数	279	100.0	17	6.1	262	93.9
	20-29歳	13	100.0	-	-	13	100.0
	30-39歳	12	100.0	-	-	12	100.0
	40-49歳	24	100.0	-	-	24	100.0
	50-59歳	42	100.0	1	2.4	41	97.6
	60-69歳	71	100.0	5	7.0	66	93.0
	70歳以上	117	100.0	11	9.4	106	90.6

注) 妊婦除外

第21表の1 血圧の状況（20歳以上、性・年齢階級別）

	総数	高血圧学会(2回測定値の平均値)														(再掲)	
		至適血圧		正常血圧		正常高値血圧		I度高血圧		II度高血圧		III度高血圧		収縮期高血圧			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%※		
総数	総数	440	100.0	89	20.2	71	16.1	112	25.5	132	30.0	29	6.6	7	1.6	104	23.6
	20-29歳	19	100.0	13	68.4	2	10.5	3	15.8	1	5.3	-	-	-	-	-	-
	30-39歳	23	100.0	12	52.2	8	34.8	2	8.7	1	4.3	-	-	-	-	-	-
	40-49歳	37	100.0	18	48.6	6	16.2	8	21.6	4	10.8	1	2.7	-	-	3	8.1
	50-59歳	64	100.0	12	18.8	15	23.4	17	26.6	14	21.9	5	7.8	1	1.6	4	6.3
	60-69歳	115	100.0	13	11.3	13	11.3	35	30.4	43	37.4	8	7.0	3	2.6	28	24.3
	70歳以上	182	100.0	21	11.5	27	14.8	47	25.8	69	37.9	15	8.2	3	1.6	69	37.9
男性	総数	159	100.0	17	10.7	27	17.0	50	31.4	45	28.3	16	10.1	4	2.5	37	23.3
	20-29歳	6	100.0	2	33.3	1	16.7	2	33.3	1	16.7	-	-	-	-	-	-
	30-39歳	10	100.0	2	20.0	6	60.0	1	10.0	1	10.0	-	-	-	-	-	-
	40-49歳	13	100.0	4	30.8	2	15.4	4	30.8	2	15.4	1	7.7	-	-	1	7.7
	50-59歳	22	100.0	-	-	5	22.7	8	36.4	4	18.2	5	22.7	-	-	1	4.5
	60-69歳	44	100.0	3	6.8	7	15.9	14	31.8	12	27.3	5	11.4	3	6.8	9	20.5
	70歳以上	64	100.0	6	9.4	6	9.4	21	32.8	25	39.1	5	7.8	1	1.6	26	40.6
女性	総数	281	100.0	72	25.6	44	15.7	62	22.1	87	31.0	13	4.6	3	1.1	67	23.8
	20-29歳	13	100.0	11	84.6	1	7.7	1	7.7	-	-	-	-	-	-	-	-
	30-39歳	13	100.0	10	76.9	2	15.4	1	7.7	-	-	-	-	-	-	-	-
	40-49歳	24	100.0	14	58.3	4	16.7	4	16.7	2	8.3	-	-	-	-	2	8.3
	50-59歳	42	100.0	12	28.6	10	23.8	9	21.4	10	23.8	-	-	1	2.4	3	7.1
	60-69歳	71	100.0	10	14.1	6	8.5	21	29.6	31	43.7	3	4.2	-	-	19	26.8
	70歳以上	118	100.0	15	12.7	21	17.8	26	22.0	44	37.3	10	8.5	2	1.7	43	36.4

注) 日本高血圧学会による血圧の分類は p.5 参照

注) 2回測定値の平均値の分布。なお、1回しか測定できなかった者については、その値を採用。

※ 総数に対する割合

第21表の2 血圧の状況（20歳以上、服薬者除外、性・年齢階級別）

	総数	高血圧学会(2回測定値の平均値)														(再掲)	
		至適血圧		正常血圧		正常高値血圧		I度高血圧		II度高血圧		III度高血圧		収縮期高血圧			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%※		
総数	総数	300	100.0	79	26.3	50	16.7	70	23.3	84	28.0	11	3.7	6	2.0	58	19.3
	20-29歳	19	100.0	13	68.4	2	10.5	3	15.8	1	5.3	-	-	-	-	-	-
	30-39歳	22	100.0	11	50.0	8	36.4	2	9.1	1	4.5	-	-	-	-	-	-
	40-49歳	37	100.0	18	48.6	6	16.2	8	21.6	4	10.8	1	2.7	-	-	3	8.1
	50-59歳	52	100.0	12	23.1	12	23.1	13	25.0	13	25.0	1	1.9	1	1.9	3	5.8
	60-69歳	81	100.0	12	14.8	5	6.2	22	27.2	34	42.0	6	7.4	2	2.5	23	28.4
	70歳以上	89	100.0	13	14.6	17	19.1	22	24.7	31	34.8	3	3.4	3	3.4	29	32.6
男性	総数	100	100.0	13	13.0	19	19.0	32	32.0	28	28.0	5	5.0	3	3.0	18	18.0
	20-29歳	6	100.0	2	33.3	1	16.7	2	33.3	1	16.7	-	-	-	-	-	-
	30-39歳	10	100.0	2	20.0	6	60.0	1	10.0	1	10.0	-	-	-	-	-	-
	40-49歳	13	100.0	4	30.8	2	15.4	4	30.8	2	15.4	1	7.7	-	-	1	7.7
	50-59歳	15	100.0	-	-	4	26.7	6	40.0	4	26.7	1	6.7	-	-	1	6.7
	60-69歳	28	100.0	2	7.1	3	10.7	9	32.1	9	32.1	3	10.7	2	7.1	6	21.4
	70歳以上	28	100.0	3	10.7	3	10.7	10	35.7	11	39.3	-	-	1	3.6	10	35.7
女性	総数	200	100.0	66	33.0	31	15.5	38	19.0	56	28.0	6	3.0	3	1.5	40	20.0
	20-29歳	13	100.0	11	84.6	1	7.7	1	7.7	-	-	-	-	-	-	-	-
	30-39歳	12	100.0	9	75.0	2	16.7	1	8.3	-	-	-	-	-	-	-	-
	40-49歳	24	100.0	14	58.3	4	16.7	4	16.7	2	8.3	-	-	-	-	2	8.3
	50-59歳	37	100.0	12	32.4	8	21.6	7	18.9	9	24.3	-	-	1	2.7	2	5.4
	60-69歳	53	100.0	10	18.9	2	3.8	13	24.5	25	47.2	3	5.7	-	-	17	32.1
	70歳以上	61	100.0	10	16.4	14	23.0	12	19.7	20	32.8	3	4.9	2	3.3	19	31.1

注) 妊婦、血圧を下げる薬服用者と服用不明者除外

注) 2回測定値の平均値の分布。なお、1回しか測定できなかった者については、その値を採用。

※ 総数に対する割合

第22表の1 収縮期（最高）・拡張期（最低）血圧の分布（20歳以上、性・年齢階級別）

2回測定値の平均値(mmHg)		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
男	収縮期（最高） 血圧	総数	159	100.0	6	100.0	10	100.0	13	100.0	22	100.0	44	100.0	64	100.0
		90未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		90~99	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		100~109	5	3.1	2	33.3	-	-	1	7.7	-	-	1	2.3	1	1.6
		110~119	12	7.5	-	-	2	20.0	3	23.1	-	-	2	4.5	5	7.8
		120~129	37	23.3	1	16.7	6	60.0	3	23.1	7	31.8	8	18.2	12	18.8
		130~139	45	28.3	2	33.3	2	20.0	4	30.8	7	31.8	15	34.1	15	23.4
		140~149	30	18.9	1	16.7	-	-	1	7.7	4	18.2	7	15.9	17	26.6
		150~159	17	10.7	-	-	-	-	1	7.7	3	13.6	5	11.4	8	12.5
		160~169	6	3.8	-	-	-	-	-	-	1	4.5	1	2.3	4	6.3
	170~179	4	2.5	-	-	-	-	-	-	-	-	3	6.8	1	1.6	
	180以上	3	1.9	-	-	-	-	-	-	-	-	2	4.5	1	1.6	
	拡張期（最低） 血圧	総数	159	100.0	6	100.0	10	100.0	13	100.0	22	100.0	44	100.0	64	100.0
		40未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		40-49	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		50-59	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		60-69	22	13.8	1	16.7	-	-	1	7.7	1	4.5	5	11.4	14	21.9
		70-79	50	31.4	1	16.7	6	60.0	5	38.5	6	27.3	11	25.0	21	32.8
		80-89	59	37.1	3	50.0	3	30.0	5	38.5	7	31.8	17	38.6	24	37.5
90-99		18	11.3	1	16.7	1	10.0	1	7.7	4	18.2	6	13.6	5	7.8	
100-109		8	5.0	-	-	-	-	1	7.7	4	18.2	3	6.8	-	-	
110-119		1	0.6	-	-	-	-	-	-	-	-	1	2.3	-	-	
120-129	1	0.6	-	-	-	-	-	-	-	-	1	2.3	-	-		
130-139	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
140以上	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
女	収縮期（最高） 血圧	総数	281	100.0	13	100.0	13	100.0	24	100.0	42	100.0	71	100.0	118	100.0
		90未満	1	0.4	1	7.7	-	-	-	-	-	-	-	-	-	-
		90~99	6	2.1	2	15.4	-	-	1	4.2	1	2.4	1	1.4	1	0.8
		100~109	24	8.5	7	53.8	6	46.2	6	25.0	1	2.4	3	4.2	1	0.8
		110~119	43	15.3	1	7.7	5	38.5	7	29.2	11	26.2	6	8.5	13	11.0
		120~129	51	18.1	1	7.7	1	7.7	6	25.0	10	23.8	10	14.1	23	19.5
		130~139	68	24.2	1	7.7	1	7.7	2	8.3	11	26.2	25	35.2	28	23.7
		140~149	50	17.8	-	-	-	-	2	8.3	4	9.5	18	25.4	26	22.0
		150~159	23	8.2	-	-	-	-	-	-	3	7.1	6	8.5	14	11.9
		160~169	11	3.9	-	-	-	-	-	-	-	-	2	2.8	9	7.6
	170~179	2	0.7	-	-	-	-	-	-	-	-	-	-	2	1.7	
	180以上	2	0.7	-	-	-	-	-	-	1	2.4	-	-	1	0.8	
	拡張期（最低） 血圧	総数	281	100.0	13	100.0	13	100.0	24	100.0	42	100.0	71	100.0	118	100.0
		40未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		40-49	1	0.4	-	-	-	-	1	4.2	-	-	-	-	-	-
		50-59	5	1.8	2	15.4	-	-	2	8.3	-	-	-	-	1	0.8
		60-69	49	17.4	5	38.5	4	30.8	3	12.5	3	7.1	11	15.5	23	19.5
		70-79	106	37.7	5	38.5	6	46.2	11	45.8	18	42.9	25	35.2	41	34.7
		80-89	84	29.9	1	7.7	3	23.1	7	29.2	13	31.0	20	28.2	40	33.9
90-99		31	11.0	-	-	-	-	-	-	7	16.7	13	18.3	11	9.3	
100-109		3	1.1	-	-	-	-	-	-	-	-	2	2.8	1	0.8	
110-119		2	0.7	-	-	-	-	-	-	1	2.4	-	-	1	0.8	
120-129	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
130-139	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
140以上	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

注) 2回測定値の平均値の分布。なお、1回しか測定できなかった者については、その値を採用。

第22表の2 収縮期（最高）・拡張期（最低）血圧の平均値、標準偏差（20歳以上、性・年齢階級別）

		総数			20-29歳			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差
男性	収縮期(最高)	159	136.8	15.7	6	123.0	13.0	10	124.7	6.1	13	127.4	13.3	22	138.0	11.9	44	140.5	18.1	64	138.8	14.5
	拡張期(最低)	159	81.3	10.7	6	81.0	11.2	10	80.1	5.2	13	81.5	11.3	22	86.1	10.8	44	84.0	12.4	64	77.9	8.6
女性	収縮期(最高)	281	131.2	17.0	13	106.9	11.7	13	112.1	8.8	24	118.0	12.1	42	128.3	15.2	71	134.7	13.3	118	137.5	16.1
	拡張期(最低)	281	77.7	10.3	13	67.5	7.9	13	73.4	6.2	24	72.8	9.7	42	80.4	10.1	71	80.0	10.3	118	78.0	10.0

注) 2回測定値の平均値の分布。なお、1回しか測定できなかった者については、その値を採用。

第22表の3 収縮期（最高）・拡張期（最低）血圧の分布（20歳以上、服薬者除外、性・年齢階級別）

2回測定値の平均値(mmHg)		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
男	収縮期（最高） 血圧	総数	100	100.0	6	100.0	10	100.0	13	100.0	15	100.0	28	100.0	28	100.0
		90未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		90~99	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		100~109	5	5.0	2	33.3	-	-	1	7.7	-	-	1	3.6	1	3.6
		110~119	8	8.0	-	-	2	20.0	3	23.1	-	-	1	3.6	2	7.1
		120~129	27	27.0	1	16.7	6	60.0	3	23.1	6	40.0	4	14.3	7	25.0
		130~139	28	28.0	2	33.3	2	20.0	4	30.8	5	33.3	9	32.1	6	21.4
		140~149	18	18.0	1	16.7	-	-	1	7.7	4	26.7	6	21.4	6	21.4
		150~159	9	9.0	-	-	-	-	1	7.7	-	-	3	10.7	5	17.9
		160~169	1	1.0	-	-	-	-	-	-	-	-	1	3.6	-	-
	170~179	2	2.0	-	-	-	-	-	-	-	-	2	7.1	-	-	
	180以上	2	2.0	-	-	-	-	-	-	-	-	1	3.6	1	3.6	
	拡張期（最低） 血圧	総数	100	100.0	6	100.0	10	100.0	13	100.0	15	100.0	28	100.0	28	100.0
		40未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		40-49	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		50-59	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		60-69	12	12.0	1	16.7	-	-	1	7.7	-	-	3	10.7	7	25.0
		70-79	31	31.0	1	16.7	6	60.0	5	38.5	6	40.0	5	17.9	8	28.6
		80-89	39	39.0	3	50.0	3	30.0	5	38.5	5	33.3	12	42.9	11	39.3
90-99		13	13.0	1	16.7	1	10.0	1	7.7	3	20.0	5	17.9	2	7.1	
100-109		4	4.0	-	-	-	-	1	7.7	1	6.7	2	7.1	-	-	
110-119		-	-	-	-	-	-	-	-	-	-	-	-	-	-	
120-129	1	1.0	-	-	-	-	-	-	-	-	1	3.6	-	-		
130-139	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
140以上	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
女	収縮期（最高） 血圧	総数	200	100.0	13	100.0	12	100.0	24	100.0	37	100.0	53	100.0	61	100.0
		90未満	1	0.5	1	7.7	-	-	-	-	-	-	-	-	-	-
		90~99	6	3.0	2	15.4	-	-	1	4.2	1	2.7	1	1.9	1	1.6
		100~109	23	11.5	7	53.8	5	41.7	6	25.0	1	2.7	3	5.7	1	1.6
		110~119	38	19.0	1	7.7	5	41.7	7	29.2	11	29.7	6	11.3	8	13.1
		120~129	35	17.5	1	7.7	1	8.3	6	25.0	8	21.6	5	9.4	14	23.0
		130~139	41	20.5	1	7.7	1	8.3	2	8.3	9	24.3	15	28.3	13	21.3
		140~149	33	16.5	-	-	-	-	2	8.3	3	8.1	15	28.3	13	21.3
		150~159	15	7.5	-	-	-	-	-	-	3	8.1	6	11.3	6	9.8
		160~169	6	3.0	-	-	-	-	-	-	-	-	2	3.8	4	6.6
	170~179	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	180以上	2	1.0	-	-	-	-	-	-	1	2.7	-	-	1	1.6	
	拡張期（最低） 血圧	総数	200	100.0	13	100.0	12	100.0	24	100.0	37	100.0	53	100.0	61	100.0
		40未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-
		40-49	1	0.5	-	-	-	-	1	4.2	-	-	-	-	-	-
		50-59	5	2.5	2	15.4	-	-	2	8.3	-	-	-	-	1	1.6
		60-69	35	17.5	5	38.5	4	33.3	3	12.5	3	8.1	9	17.0	11	18.0
		70-79	79	39.5	5	38.5	5	41.7	11	45.8	17	45.9	19	35.8	22	36.1
		80-89	55	27.5	1	7.7	3	25.0	7	29.2	9	24.3	14	26.4	21	34.4
90-99		20	10.0	-	-	-	-	-	-	7	18.9	9	17.0	4	6.6	
100-109		3	1.5	-	-	-	-	-	-	-	-	2	3.8	1	1.6	
110-119		2	1.0	-	-	-	-	-	-	1	2.7	-	-	1	1.6	
120-129	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
130-139	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
140以上	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

注) 妊婦、血圧を下げる薬服用者と服用不明者除外。

注) 2回測定値の平均値の分布。なお、1回しか測定できなかった者については、その値を採用。

第22表の4 収縮期（最高）・拡張期（最低）血圧の平均値、標準偏差（20歳以上、服薬者除外、性・年齢階級別）

(mmHg)

		総数			20-29歳			30-39歳			40-49歳			50-59歳			60-69歳			70歳以上		
		人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差
男性	収縮期(最高)	100	134.3	15.0	6	123.0	13.0	10	124.7	6.1	13	127.4	13.3	15	133.7	8.4	28	140.8	16.8	28	137.2	14.9
	拡張期(最低)	100	81.9	10.8	6	81.0	11.2	10	80.1	5.2	13	81.5	11.3	15	84.4	9.2	28	85.5	12.4	28	78.0	9.2
女性	収縮期(最高)	200	128.5	17.4	13	106.9	11.7	12	112.8	8.8	24	118.0	12.1	37	128.1	15.8	53	135.0	15.0	61	134.9	16.5
	拡張期(最低)	200	77.3	10.8	13	67.5	7.9	12	73.7	6.4	24	72.8	9.7	37	80.3	10.6	53	79.5	10.7	61	78.0	10.7

注) 妊婦、血圧を下げる薬服用者と服用不明者除外。

注) 2回測定値の平均値の分布。なお、1回しか測定できなかった者については、その値を採用。

第23表の1 ヘモグロビンA1cの分布（20歳以上、総数・年齢階級別）

総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)											
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	65-69歳		70-74歳		75-79歳		80-84歳		85歳以上			
													人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	380	100.0																						
3.9%以下	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.0-4.1%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.2-4.3%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.4-4.5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.6-4.7%	1	0.3	-	-	-	-	-	-	-	-	-	1	0.7	-	-	-	-	-	-	-	-	1	5.6	-
4.8-4.9%	5	1.3	-	-	2	9.5	1	2.8	1	1.0	1	0.7	-	-	-	-	-	-	-	-	-	-	-	-
5.0-5.1%	22	5.8	5	33.3	5	23.8	2	5.6	1	1.0	4	2.7	1	2.3	1	2.3	2	3.8	1	3.0	1	3.0	-	-
5.2%	19	5.0	4	26.7	2	9.5	4	11.1	3	4.0	2	1.4	2	3.3	-	-	1	1.9	-	-	1	5.6	-	
5.3%	19	5.0	1	6.7	-	-	5	13.9	3	5.0	5	3.4	2	3.3	1	2.3	2	3.8	1	3.0	1	5.6	-	
5.4%	29	7.6	1	6.7	5	23.8	6	16.7	4	6.8	2	2.0	11	7.4	3	6.8	3	5.7	5	15.2	-	-	-	
5.5%	31	8.2	-	-	1	4.8	5	13.9	6	10.2	8	7.9	11	7.4	4	6.6	4	7.5	1	3.0	3	16.7	-	
5.6%	31	8.2	1	6.7	2	9.5	6	16.7	6	10.2	11	10.9	5	3.4	7	11.5	2	4.5	-	-	2	6.1	1	
5.7%	44	11.6	2	13.3	2	9.5	1	2.8	10	16.9	14	13.9	15	10.1	10	16.4	6	13.6	6	13.6	2	6.1	1	
5.8%	32	8.4	1	6.7	2	9.5	1	2.8	3	3.4	10	9.9	16	10.8	7	11.5	6	13.2	7	13.2	3	9.1	-	
5.9%	31	8.2	-	-	-	-	1	2.8	6	10.2	10	9.9	14	9.5	14	9.5	4	9.1	4	7.5	3	9.1	3	
6.0%	26	6.8	-	-	-	-	1	2.8	7	11.9	8	7.9	10	6.8	8	9.8	4	9.4	5	9.4	4	12.1	1	
6.1%	23	6.1	-	-	-	-	2	5.6	4	6.8	6	5.9	11	7.4	4	6.6	5	11.4	3	5.7	3	9.1	-	
6.2-6.3%	23	6.1	-	-	-	-	1	2.8	1	1.7	10	9.9	11	7.4	7	11.5	4	9.1	5	9.4	1	3.0	1	
6.4-6.5%	5	1.3	-	-	-	-	-	-	1	1.0	1	1.0	4	2.7	1	1.6	1	2.3	1	1.9	2	6.1	-	
6.6-6.7%	14	3.7	-	-	-	-	1	1.7	1	1.7	2	2.0	11	7.4	1	1.6	3	6.8	3	5.7	3	9.1	2	
6.8-6.9%	8	2.1	-	-	-	-	-	-	4	4.0	4	4.0	4	2.7	3	4.9	1	2.3	2	3.8	-	1	5.6	
7.0-7.1%	3	0.8	-	-	-	-	-	-	-	-	3	2.0	3	2.0	-	-	1	1.9	2	6.1	-	-	-	
7.2-7.3%	3	0.8	-	-	-	-	-	-	1	1.0	2	1.4	2	1.4	1	1.6	1	2.3	-	-	-	1	5.6	
7.4-7.5%	1	0.3	-	-	-	-	-	-	-	-	1	0.7	1	0.7	-	-	1	2.3	-	-	-	-	-	
7.6-7.7%	4	1.1	-	-	-	-	-	-	2	2.0	2	1.4	2	1.4	1	1.6	1	2.3	1	1.9	-	-	-	
7.8-7.9%	1	0.3	-	-	-	-	-	-	-	-	1	0.7	1	0.7	-	-	1	1.9	-	-	-	-	-	
8.0-8.1%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.2-8.3%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.4-8.5%	2	0.5	-	-	-	-	-	-	-	-	2	1.4	2	1.4	-	-	1	1.9	-	-	-	1	5.6	
8.6-8.7%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.8-8.9%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
9.0%以上	3	0.8	-	-	-	-	-	-	1	1.7	1	1.0	1	0.7	1	1.6	-	-	1	1.9	-	-	-	
(再掲) 6.5%以上	40	10.5	-	-	-	-	-	-	2	3.4	10	9.9	28	18.9	7	11.5	8	18.2	10	18.9	5	15.2	5	27.8

第23表の2 ヘモグロビンA1cの分布（20歳以上、男性・年齢階級別）

総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)										
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	65-69歳		70-74歳		75-79歳		80-84歳		85歳以上		
													人数	%	人数	%	人数	%	人数	%	人数	%	人数
総数	132	100.0	-	-	13	100.0	21	100.0	35	100.0	49	100.0	23	100.0	13	100.0	17	100.0	13	100.0	6	100.0	
3.9%以下	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.0-4.1%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.2-4.3%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.4-4.5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.6-4.7%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.8-4.9%	3	2.3	-	-	1	11.1	-	-	1	2.9	-	-	-	-	-	-	-	-	-	-	-	-	-
5.0-5.1%	11	8.3	2	40.0	3	33.3	3	14.3	1	2.9	1	2.0	1	4.3	1	4.3	1	5.9	1	7.7	1	16.7	
5.2%	5	3.8	1	20.0	1	11.1	-	-	1	2.9	2	4.1	-	-	-	-	-	-	-	-	-	-	
5.3%	9	6.8	1	20.0	-	-	2	9.5	2	5.7	1	2.0	1	4.3	1	4.3	1	7.7	-	-	-	-	
5.4%	6	4.5	-	-	1	11.1	2	15.4	-	-	1	2.0	-	-	1	7.7	-	-	-	-	-	-	
5.5%	9	6.8	-	-	-	-	3	14.3	3	8.6	2	4.1	1	4.3	1	7.7	-	-	-	-	-	-	
5.6%	14	10.6	-	-	2	22.2	3	23.1	3	8.6	1	2.0	3	13.0	-	-	-	-	-	-	1	16.7	
5.7%	11	8.3	1	20.0	-	-	2	9.5	3	8.6	5	10.2	2	8.7	1	7.7	3	17.6	1	7.7	-	-	
5.8%	11	8.3	-	-	1	11.1	1	7.7	4	11.4	5	10.2	2	8.7	2	15.4	2	11.8	1	7.7	-	-	
5.9%	10	7.6	-	-	-	-	-	-	2	5.7	7	14.3	2	8.7	2	15.4	1	5.9	3	23.1	1	16.7	
6.0%	9	6.8	-	-	-	-	-	-	3	8.6	6	12.2	1	4.3	-	-	3	17.6	3	23.1	-	-	
6.1%	10	7.6	-	-	-	-	-	-	3	8.6	5	10.2	2	8.7	3	23.1	-	-	2	15.4	-	-	
6.2-6.3%	6	4.5	-	-	-	-	-	-	4	11.4	2	4.1	3	13.0	1	7.7	1	5.9	-	-	-	-	
6.4-6.5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
6.6-6.7%	3	2.3	-	-	-	-	1	4.8	-	-	2	4.1	-	-	-	-	-	-	-	-	-	-	
6.8-6.9%	3	2.3	-	-	-	-	-	-	2	5.7	1	2.0	2	8.7	-	-	1	5.9	-	-	-	-	
7.0-7.1%	2	1.5	-	-	-	-	-	-	-	-	2	4.1	-	-	-	-	1	5.9	-	-	-	-	
7.2-7.3%	2	1.5	-	-	-	-	-	-	1	2.9	1	2.0	1	4.3	1	7.7	-	-	-	-	-	-	
7.4-7.5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
7.6-7.7%	2	1.5	-	-	-	-	-	-	1	2.9	1	2.0	1	4.3	-	-	1	5.9	-	-	-	-	
7.8-7.9%	1	0.8	-	-	-	-	-	-	-	-	1	2.0	-	-	-	-	1	5.9	-	-	-	-	
8.0-8.1%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.2-8.3%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.4-8.5%	2	1.5	-	-	-	-	-	-	-	-	2	4.1	-	-	-	-	1	5.9	-	-	1	16.7	
8.6-8.7%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.8-8.9%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
9.0%以上	3	2.3	-	-	-	-	1	4.8	1	2.9	1	2.0	1	4.3	-	-	1	5.9	-	-	-	-	
(再掲) 6.5%以上	18	13.6	-	-	-	-	2	9.5	5	14.3	11	22.4	5	21.7	1	7.7	6	35.3	2	15.4	2	33.3	

第23表の3 ヘモグロビンA1cの分布（20歳以上、女性・年齢階級別）

年齢階級	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)											
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	65-69歳		70-74歳		75-79歳		80-84歳		85歳以上			
															人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
3.9%以下	248	100.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.0-4.1%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.2-4.3%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.4-4.5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.6-4.7%	1	0.4	-	-	-	-	-	-	-	-	-	-	-	1	1.0	-	-	-	-	-	-	-	-	1	8.3	-
4.8-4.9%	2	0.8	-	-	1	8.3	-	-	-	-	-	-	1	1.0	-	-	-	-	-	-	-	-	-	-	-	-
5.0-5.1%	11	4.4	3	30.0	2	16.7	1	4.3	2	5.3	-	-	3	3.0	-	-	1	3.2	2	5.6	-	-	-	-	-	-
5.2%	14	5.6	3	30.0	1	8.3	4	17.4	3	7.9	3	4.5	-	-	-	-	2	5.3	-	-	-	-	-	-	-	-
5.3%	10	4.0	-	-	-	-	2	8.7	1	2.6	3	4.5	4	4.0	-	-	1	2.6	-	-	2	5.6	1	5.0	1	8.3
5.4%	23	9.3	1	10.0	4	33.3	4	17.4	2	5.3	2	3.0	10	10.1	-	-	2	6.5	3	8.3	3	5	25.0	-	-	
5.5%	22	8.9	-	-	1	8.3	4	17.4	3	7.9	5	7.6	9	9.1	-	-	2	6.5	4	11.1	4	1	5.0	2	16.7	
5.6%	17	6.9	1	10.0	-	-	3	13.0	1	2.6	8	12.1	4	4.0	-	-	2	6.5	2	10.0	-	-	-	-	-	
5.7%	33	13.3	1	10.0	2	16.7	1	4.3	8	21.1	11	16.7	10	10.1	-	-	5	16.1	3	8.3	5	2	10.0	1	8.3	
5.8%	21	8.5	1	10.0	1	8.3	-	-	2	5.3	6	9.1	11	11.1	-	-	4	12.9	5	13.9	2	1	5.0	-	-	
5.9%	21	8.5	-	-	-	-	-	-	6	15.8	8	12.1	7	7.1	-	-	2	6.5	3	8.3	3	2	10.0	2	16.7	
6.0%	17	6.9	-	-	-	-	1	4.3	7	18.4	5	7.6	4	4.0	-	-	4	10.5	2	5.6	2	1	5.0	1	8.3	
6.1%	13	5.2	-	-	-	-	2	8.7	2	5.3	3	4.5	6	6.1	-	-	2	6.5	3	8.3	3	1	5.0	-	-	
6.2-6.3%	17	6.9	-	-	-	-	1	4.3	1	2.6	6	9.1	9	9.1	-	-	4	10.5	3	11.1	4	1	5.0	1	8.3	
6.4-6.5%	5	2.0	-	-	-	-	-	-	-	-	1	1.5	4	4.0	-	-	1	3.2	1	2.8	1	2	10.0	-	-	
6.6-6.7%	11	4.4	-	-	-	-	-	-	-	-	2	3.0	9	9.1	-	-	3	9.7	3	8.3	3	2	10.0	1	8.3	
6.8-6.9%	5	2.0	-	-	-	-	-	-	-	-	2	3.0	3	3.0	-	-	1	3.2	1	2.8	1	1	5.0	1	8.3	
7.0-7.1%	1	0.4	-	-	-	-	-	-	-	-	-	-	1	1.0	-	-	-	-	-	-	-	1	5.0	-	-	
7.2-7.3%	1	0.4	-	-	-	-	-	-	-	-	-	-	1	1.0	-	-	-	-	-	-	-	-	-	1	8.3	
7.4-7.5%	1	0.4	-	-	-	-	-	-	-	-	-	-	1	1.0	-	-	1	3.2	-	-	-	-	-	-	-	
7.6-7.7%	2	0.8	-	-	-	-	-	-	-	-	1	1.5	1	1.0	-	-	1	3.2	-	-	-	-	-	-	-	
7.8-7.9%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.0-8.1%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.2-8.3%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.4-8.5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.6-8.7%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
8.8-8.9%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
9.0%以上	22	8.9	-	-	-	-	-	-	-	-	5	7.6	17	17.2	-	-	2	22.6	4	11.1	3	3	15.0	3	25.0	
(再掲) 6.5%以上	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

第23表の4 ヘモグロビンA1cの分布（20歳以上、服薬者除外、総数・年齢階級別）

総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)											
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	65-69歳		70-74歳		75-79歳		80-84歳		85歳以上			
													人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	360	100.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
3.9%以下	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.0-4.1%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.2-4.3%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.4-4.5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.6-4.7%	1	0.3	-	-	-	-	-	-	-	-	-	1	0.7	-	-	-	-	-	-	-	-	1	6.7	-
4.8-4.9%	5	1.4	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5.0-5.1%	22	6.1	5	33.3	2	5.6	5	8.5	1	1.1	1	1.1	1	2.8	1	2.4	2	2.4	1	1.8	1	3.2	-	-
5.2%	18	5.0	4	26.7	1	5.0	4	11.1	3	5.1	4	4.2	2	3.5	2	3.5	2	2.4	1	2.1	-	1	6.7	-
5.3%	19	5.3	1	6.7	-	-	5	13.9	3	5.1	5	5.3	5	3.7	2	3.5	2	2.4	2	4.2	1	3.2	1	6.7
5.4%	29	8.1	1	6.7	5	25.0	6	16.7	4	6.8	2	2.1	11	8.1	-	-	3	7.3	3	6.3	5	16.1	-	-
5.5%	31	8.6	-	-	1	5.0	5	13.9	6	10.2	8	8.4	11	8.1	4	7.0	3	7.3	4	8.3	1	3.2	3	20.0
5.6%	31	8.6	1	6.7	2	10.0	6	16.7	6	10.2	11	11.6	5	3.7	7	12.3	2	4.9	-	2	2	6.5	1	6.7
5.7%	44	12.2	2	13.3	2	10.0	1	2.8	10	16.9	14	14.7	15	11.1	10	17.5	6	14.6	6	12.5	2	6.5	1	6.7
5.8%	32	8.9	1	6.7	2	10.0	1	2.8	2	3.4	10	10.5	16	11.9	7	12.3	6	14.6	7	14.6	3	9.7	-	-
5.9%	30	8.3	-	-	-	-	1	2.8	6	10.2	10	10.5	13	9.6	6	10.5	4	9.8	3	6.3	3	9.7	3	20.0
6.0%	26	7.2	-	-	-	-	1	2.8	7	11.9	8	8.4	10	7.4	3	5.3	-	-	5	10.4	4	12.9	1	6.7
6.1%	22	6.1	-	-	-	-	2	5.6	4	6.8	6	6.3	10	7.4	4	7.0	5	12.2	3	6.3	2	6.5	-	-
6.2-6.3%	20	5.6	-	-	-	-	1	2.8	1	1.7	8	8.4	10	7.4	6	10.5	3	7.3	5	10.4	1	3.2	1	6.7
6.4-6.5%	5	1.4	-	-	-	-	-	-	-	-	1	1.1	4	3.0	1	1.8	1	2.4	1	2.1	2	6.5	-	-
6.6-6.7%	11	3.1	-	-	-	-	-	-	1	1.7	1	1.1	9	6.7	-	-	2	4.9	3	6.3	3	9.7	1	6.7
6.8-6.9%	6	1.7	-	-	-	-	-	-	-	-	4	4.2	2	1.5	3	5.3	1	2.4	1	2.1	-	-	-	-
7.0-7.1%	1	0.3	-	-	-	-	-	-	-	-	-	-	1	0.7	-	-	-	-	-	-	1	3.2	-	-
7.2-7.3%	2	0.6	-	-	-	-	-	-	-	-	-	-	1	0.7	1	1.8	1	2.4	-	-	-	-	-	-
7.4-7.5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
7.6-7.7%	1	0.3	-	-	-	-	-	-	-	-	-	-	1	0.7	-	-	-	2.4	-	-	-	-	-	-
7.8-7.9%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8.0-8.1%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8.2-8.3%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8.4-8.5%	2	0.6	-	-	-	-	-	-	-	-	-	-	2	1.5	-	-	-	-	-	-	-	-	-	-
8.6-8.7%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8.8-8.9%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
9.0%以上	2	0.6	-	-	-	-	1	1.7	-	-	-	-	1	0.7	-	-	-	-	-	-	-	-	-	-
(再掲)6.5%以上	26	7.2	-	-	-	-	2	3.4	6	6.3	18	13.3	4	7.0	6	14.6	6	14.6	6	12.5	4	12.9	2	13.3

(注) インスリン注射または血糖を下げる薬服用者と服用不明者除外

第23表の5 ヘモグロビンA1cの分布（20歳以上、服薬者除外、男性・年齢階級別）

総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)											
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	65-69歳		70-74歳		75-79歳		80-84歳		85歳以上			
													人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	123	100.0	5	100.0	9	100.0	13	100.0	21	100.0	32	100.0	43	100.0	20	100.0	13	100.0	12	100.0	12	100.0	6	100.0
3.9%以下	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.0-4.1%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.2-4.3%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.4-4.5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.6-4.7%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
4.8-4.9%	3	2.4	-	-	1	11.1	1	7.7	-	-	1	3.1	-	-	-	-	-	-	-	-	-	-	-	-
5.0-5.1%	11	8.9	2	40.0	3	33.3	1	7.7	3	14.3	1	3.1	1	2.3	1	5.0	-	-	1	8.3	-	-	-	-
5.2%	5	4.1	1	20.0	1	11.1	-	-	-	-	1	3.1	2	4.7	-	-	1	8.3	-	-	-	1	16.7	
5.3%	9	7.3	1	20.0	-	-	3	23.1	2	9.5	2	6.3	1	2.3	1	5.0	-	-	-	-	-	-	-	-
5.4%	6	4.9	-	-	1	11.1	2	15.4	2	9.5	-	-	1	2.3	-	-	-	-	-	-	-	-	-	-
5.5%	9	7.3	-	-	-	-	1	7.7	3	14.3	3	9.4	2	4.7	1	5.0	-	-	-	-	-	-	1	16.7
5.6%	14	11.4	-	-	2	22.2	3	23.1	5	23.8	3	9.4	1	2.3	3	15.0	-	-	-	-	-	-	1	16.7
5.7%	11	8.9	1	20.0	-	-	-	-	2	9.5	3	9.4	5	11.6	2	10.0	1	7.7	3	25.0	1	8.3	-	-
5.8%	11	8.9	-	-	1	11.1	1	7.7	-	-	4	12.5	5	11.6	2	10.0	2	15.4	2	16.7	1	8.3	-	-
5.9%	9	7.3	-	-	-	-	1	7.7	-	-	2	6.3	6	14.0	2	10.0	2	15.4	-	-	3	25.0	1	16.7
6.0%	9	7.3	-	-	-	-	-	-	-	-	3	9.4	6	14.0	1	5.0	-	-	3	25.0	3	25.0	-	-
6.1%	9	7.3	-	-	-	-	-	2	9.5	3	9.4	4	9.3	2	10.0	3	23.1	-	-	1	8.3	-	-	
6.2-6.3%	5	4.1	-	-	-	-	-	-	-	-	3	9.4	2	4.7	2	10.0	1	7.7	1	8.3	-	-	-	-
6.4-6.5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
6.6-6.7%	3	2.4	-	-	1	4.8	-	-	1	4.8	-	-	2	4.7	-	-	-	-	-	-	1	8.3	1	16.7
6.8-6.9%	2	1.6	-	-	-	-	-	-	-	-	2	6.3	1	2.3	2	10.0	-	-	-	-	-	-	-	-
7.0-7.1%	1	0.8	-	-	-	-	-	-	-	-	-	-	1	2.3	-	-	-	-	-	-	1	8.3	-	-
7.2-7.3%	2	1.6	-	-	-	-	-	-	-	-	1	3.1	1	2.3	1	5.0	1	7.7	-	-	-	-	-	-
7.4-7.5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
7.6-7.7%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
7.8-7.9%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8.0-8.1%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8.2-8.3%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8.4-8.5%	2	1.6	-	-	-	-	-	-	-	-	-	-	2	4.7	-	-	-	-	-	-	-	-	1	16.7
8.6-8.7%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
8.8-8.9%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
9.0%以上	2	1.6	-	-	-	-	-	-	1	4.8	-	-	1	2.3	-	-	-	-	-	-	-	-	-	-
(再掲)6.5%以上	12	9.8	-	-	-	-	-	-	2	9.5	3	9.4	7	16.3	3	15.0	1	7.7	2	16.7	2	16.7	2	33.3

(注) インスリン注射または血糖を下げる薬服用者と服用不明者除外

第23表の6 ヘモグロビンA1cの分布（20歳以上、服薬者除外、女性・年齢階級別）

総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)											
	人数		%		人数		%		人数		%		人数		%		人数		%		人数		%	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
237	100.0	10	100.0	11	100.0	23	100.0	38	100.0	63	100.0	92	100.0	37	100.0	28	100.0	36	100.0	19	100.0	9	100.0	
総数																								
3.9%以下																								
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1	0.4	-	-	-	-	-	-	-	-	-	-	1	1.1	-	-	-	-	-	-	-	-	1	11.1	
2	0.8	-	-	1	9.1	-	-	-	-	-	-	1	1.1	-	-	1	3.6	-	-	-	-	-	-	
11	4.6	3	30.0	2	18.2	1	4.3	2	5.3	-	-	3	3.3	-	-	1	3.6	2	5.6	-	-	-	-	
13	5.5	3	30.0	-	-	4	17.4	3	7.9	3	4.8	-	-	2	5.4	-	-	-	-	-	-	-	-	
10	4.2	-	-	-	-	2	8.7	1	2.6	3	4.8	4	4.3	1	2.7	-	-	2	5.6	1	5.3	1	11.1	
23	9.7	1	10.0	4	36.4	4	17.4	2	5.3	2	3.2	10	10.9	-	-	2	7.1	3	8.3	5	26.3	2	22.2	
22	9.3	-	-	1	9.1	4	17.4	3	7.9	5	7.9	9	9.8	3	8.1	2	7.1	4	11.1	1	5.3	2	22.2	
17	7.2	1	10.0	-	-	3	13.0	1	2.6	8	12.7	4	4.3	4	10.8	2	7.1	-	-	2	10.5	-	-	
33	13.9	1	10.0	2	18.2	1	4.3	8	21.1	11	17.5	10	10.9	8	21.6	5	17.9	3	8.3	1	5.3	1	11.1	
21	8.9	1	10.0	1	9.1	-	-	2	5.3	6	9.5	11	12.0	5	13.5	4	14.3	5	13.9	2	10.5	-	-	
21	8.9	-	-	-	-	-	-	6	15.8	8	12.7	7	7.6	4	10.8	2	7.1	3	8.3	-	-	2	22.2	
17	7.2	-	-	-	-	1	4.3	7	18.4	5	7.9	4	4.3	2	5.4	-	-	2	5.6	1	5.3	1	11.1	
13	5.5	-	-	-	-	2	8.7	2	5.3	3	4.8	6	6.5	2	5.4	2	7.1	3	8.3	1	5.3	-	-	
15	6.3	-	-	-	-	1	4.3	1	2.6	5	7.9	8	8.7	4	10.8	2	7.1	4	11.1	1	5.3	1	11.1	
5	2.1	-	-	-	-	-	-	-	-	1	1.6	4	4.3	1	2.7	1	3.6	1	2.8	1	10.5	-	-	
8	3.4	-	-	-	-	-	-	-	-	1	1.6	7	7.6	-	-	2	7.1	3	8.3	2	10.5	-	-	
4	1.7	-	-	-	-	-	-	-	-	2	3.2	2	2.2	1	2.7	1	3.6	1	2.8	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
-	-	-	-	-																				

第 24 表の 1 ヘモグロビン A1c の平均値、標準偏差 (20 歳以上、性・年齢階級別)

(%)

年齢(歳)	総数			男性			女性		
	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差
総数	380	5.8	0.7	132	5.9	1.0	248	5.8	0.5
20-29歳	15	5.3	0.3	5	5.3	0.2	10	5.3	0.3
30-39歳	21	5.3	0.3	9	5.3	0.3	12	5.4	0.3
40-49歳	36	5.5	0.3	13	5.4	0.3	23	5.5	0.3
50-59歳	59	5.7	0.7	21	5.8	1.1	38	5.7	0.3
60-69歳	101	5.9	0.6	35	6.0	0.8	66	5.9	0.4
70歳以上	148	6.0	0.8	49	6.3	1.2	99	5.9	0.6
(再掲)									
65-69歳	61	6.0	0.6	23	6.2	0.9	38	5.8	0.3
70-74歳	44	6.0	0.6	13	5.9	0.5	31	6.0	0.6
75-79歳	53	6.1	1.1	17	6.7	1.7	36	5.9	0.4
80-84歳	33	5.9	0.5	13	6.0	0.4	20	5.9	0.5
85歳以上	18	6.0	0.9	6	6.2	1.1	12	6.0	0.7

第 24 表の 2 ヘモグロビン A1c の平均値、標準偏差 (20 歳以上、服薬者除外、性・年齢階級別)

(%)

年齢(歳)	総数			男性			女性		
	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差
総数	360	5.8	0.7	123	5.9	1.0	237	5.7	0.4
20-29歳	15	5.3	0.3	5	5.3	0.2	10	5.3	0.3
30-39歳	20	5.4	0.3	9	5.3	0.3	11	5.4	0.3
40-49歳	36	5.5	0.3	13	5.4	0.3	23	5.5	0.3
50-59歳	59	5.7	0.7	21	5.8	1.1	38	5.7	0.3
60-69歳	95	5.8	0.4	32	5.9	0.5	63	5.8	0.4
70歳以上	135	6.0	0.8	43	6.2	1.2	92	5.8	0.5
(再掲)									
65-69歳	57	5.9	0.4	20	6.0	0.5	37	5.8	0.3
70-74歳	41	5.9	0.5	13	5.9	0.5	28	5.9	0.6
75-79歳	48	6.1	1.1	12	6.6	2.0	36	5.9	0.4
80-84歳	31	5.9	0.5	12	6.0	0.5	19	5.8	0.4
85歳以上	15	5.9	0.8	6	6.2	1.1	9	5.6	0.5

注) インスリン注射または血糖を下げる薬服用者と服用不明者除外

第25表の1 総コレステロール値の分布（20歳以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)											
															65-69歳		70-74歳		75-79歳		80-84歳		85歳以上			
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
男性	総数	132	100.0	5	100.0	9	100.0	13	100.0	21	100.0	35	100.0	49	100.0	23	100.0	13	100.0	17	100.0	13	100.0	6	100.0	
	120mg/dl未満	1	0.8	-	-	-	-	-	-	-	-	-	-	1	2.0	-	-	-	-	1	5.9	-	-	-	-	
	120-140mg/dl未満	9	6.8	-	-	-	-	1	7.7	1	4.8	1	2.9	6	12.2	1	4.3	-	-	2	11.8	2	15.4	2	33.3	
	140-160mg/dl未満	12	9.1	1	20.0	-	-	1	7.7	1	4.8	2	5.7	7	14.3	1	4.3	2	15.4	2	11.8	2	15.4	1	16.7	
	160-180mg/dl未満	25	18.9	2	40.0	2	22.2	2	15.4	4	19.0	3	8.6	12	24.5	1	4.3	3	23.1	5	29.4	2	15.4	2	33.3	
	180-200mg/dl未満	26	19.7	1	20.0	1	11.1	2	15.4	4	19.0	10	28.6	8	16.3	6	26.1	4	30.8	3	17.6	1	7.7	-	-	
	200-220mg/dl未満	27	20.5	-	-	3	33.3	2	15.4	6	28.6	7	20.0	9	18.4	5	21.7	3	23.1	3	17.6	3	23.1	-	-	
	220-240mg/dl未満	17	12.9	-	-	3	33.3	5	38.5	2	9.5	4	11.4	3	6.1	3	13.0	-	-	1	5.9	1	7.7	1	16.7	
	240-260mg/dl未満	7	5.3	1	20.0	-	-	-	-	1	4.8	4	11.4	1	2.0	3	13.0	-	-	-	-	-	1	7.7	-	-
	260-280mg/dl未満	4	3.0	-	-	-	-	-	-	1	4.8	2	5.7	1	2.0	1	4.3	-	-	-	-	-	1	7.7	-	-
	280-300mg/dl未満	1	0.8	-	-	-	-	-	-	-	-	-	1	2.9	-	-	1	4.3	-	-	-	-	-	-	-	
300mg/dl以上	3	2.3	-	-	-	-	-	-	1	4.8	1	2.9	1	2.0	1	4.3	1	7.7	-	-	-	-	-	-		
女性	総数	248	100.0	10	100.0	12	100.0	23	100.0	38	100.0	66	100.0	99	100.0	38	100.0	31	100.0	36	100.0	20	100.0	12	100.0	
	120mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	120-140mg/dl未満	5	2.0	-	-	1	8.3	1	4.3	-	-	1	1.5	2	2.0	-	-	-	-	-	-	1	5.0	1	8.3	
	140-160mg/dl未満	12	4.8	1	10.0	1	8.3	3	13.0	4	10.5	1	1.5	2	2.0	1	2.6	1	3.2	-	-	-	-	1	8.3	
	160-180mg/dl未満	34	13.7	4	40.0	4	33.3	3	13.0	4	10.5	6	9.1	13	13.1	2	5.3	1	3.2	5	13.9	4	20.0	3	25.0	
	180-200mg/dl未満	50	20.2	2	20.0	2	16.7	7	30.4	5	13.2	9	13.6	25	25.3	7	18.4	9	29.0	11	30.6	4	20.0	1	8.3	
	200-220mg/dl未満	52	21.0	1	10.0	2	16.7	4	17.4	8	21.1	14	21.2	23	23.2	7	18.4	5	16.1	10	27.8	4	20.0	4	33.3	
	220-240mg/dl未満	44	17.7	1	10.0	1	8.3	2	8.7	9	23.7	18	27.3	13	13.1	12	31.6	6	19.4	4	11.1	2	10.0	1	8.3	
	240-260mg/dl未満	26	10.5	-	-	-	-	1	4.3	4	10.5	7	10.6	14	14.1	3	7.9	8	25.8	3	8.3	3	15.0	-	-	
	260-280mg/dl未満	14	5.6	-	-	-	-	1	4.3	2	5.3	7	10.6	4	4.0	4	10.5	1	3.2	1	2.8	2	10.0	-	-	
	280-300mg/dl未満	9	3.6	1	10.0	1	8.3	1	4.3	1	2.6	2	3.0	3	3.0	1	2.6	-	-	2	5.6	-	-	1	8.3	
300mg/dl以上	2	0.8	-	-	-	-	-	-	1	2.6	1	1.5	-	-	1	2.6	-	-	-	-	-	-	-	-		

第25表の2 総コレステロール値の平均値、標準偏差（20歳以上、性・年齢階級別）

(mg/dl)

年齢(歳)	男性			女性		
	人数	平均値	標準偏差	人数	平均値	標準偏差
総数	132	196.3	39.7	248	210.1	36.7
20-29歳	5	182.0	33.3	10	191.9	40.7
30-39歳	9	204.7	24.4	12	187.5	38.1
40-49歳	13	197.4	33.0	23	196.0	37.6
50-59歳	21	202.8	41.7	38	213.2	37.6
60-69歳	35	211.6	39.2	66	220.2	35.7
70歳以上	49	182.3	38.3	99	210.0	33.0
(再掲)						
65-69歳	23	216.5	41.4	38	221.7	34.9
70-74歳	13	192.4	39.8	31	217.1	27.9
75-79歳	17	174.4	31.9	36	208.4	30.6
80-84歳	13	190.8	41.6	20	210.4	37.1
85歳以上	6	164.7	32.6	12	195.9	39.0

第 25 表の 3 総コレステロール値の分布（20 歳以上、服薬者除外、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)											
															65-69歳		70-74歳		75-79歳		80-84歳		85歳以上			
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%		
男性	総数	111	100.0	5	100.0	9	100.0	13	100.0	17	100.0	31	100.0	36	100.0	19	100.0	10	100.0	13	100.0	7	100.0	6	100.0	
	120mg/dl未満	1	0.9	-	-	-	-	-	-	-	-	-	-	1	2.8	-	-	-	-	1	7.7	-	-	-	-	
	120-140mg/dl未満	8	7.2	-	-	-	-	1	7.7	1	5.9	1	3.2	5	13.9	1	5.3	-	-	2	15.4	1	14.3	2	33.3	
	140-160mg/dl未満	10	9.0	1	20.0	-	-	1	7.7	1	5.9	2	6.5	5	13.9	1	5.3	1	10.0	1	7.7	2	28.6	1	16.7	
	160-180mg/dl未満	19	17.1	2	40.0	2	22.2	2	15.4	4	23.5	2	6.5	7	19.4	-	-	2	20.0	2	15.4	1	14.3	2	33.3	
	180-200mg/dl未満	24	21.6	1	20.0	1	11.1	2	15.4	3	17.6	9	29.0	8	22.2	5	26.3	4	40.0	3	23.1	1	14.3	-	-	
	200-220mg/dl未満	23	20.7	-	-	3	33.3	2	15.4	5	29.4	6	19.4	7	19.4	4	21.1	3	30.0	3	23.1	1	14.3	-	-	
	220-240mg/dl未満	16	14.4	-	-	3	33.3	5	38.5	2	11.8	4	12.9	2	5.6	3	15.8	-	-	1	7.7	-	-	1	16.7	
	240-260mg/dl未満	5	4.5	1	20.0	-	-	-	-	-	-	3	9.7	1	2.8	2	10.5	-	-	-	-	-	-	1	14.3	
	260-280mg/dl未満	2	1.8	-	-	-	-	-	-	-	-	2	6.5	-	-	1	5.3	-	-	-	-	-	-	-	-	
	280-300mg/dl未満	1	0.9	-	-	-	-	-	-	-	-	1	3.2	-	-	1	5.3	-	-	-	-	-	-	-	-	
	300mg/dl以上	2	1.8	-	-	-	-	-	-	1	5.9	1	3.2	-	-	1	5.3	-	-	-	-	-	-	-	-	
女性	総数	185	100.0	10	100.0	11	100.0	23	100.0	31	100.0	42	100.0	68	100.0	26	100.0	24	100.0	23	100.0	14	100.0	7	100.0	
	120mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	120-140mg/dl未満	3	1.6	-	-	1	9.1	1	4.3	-	-	-	-	1	1.5	-	-	-	-	-	-	-	-	-	1	14.3
	140-160mg/dl未満	9	4.9	1	10.0	1	9.1	3	13.0	3	9.7	-	-	1	1.5	-	-	-	-	-	-	-	-	-	1	14.3
	160-180mg/dl未満	19	10.3	4	40.0	3	27.3	3	13.0	2	6.5	1	2.4	6	8.8	1	3.8	1	4.2	2	8.7	1	7.1	2	28.6	
	180-200mg/dl未満	33	17.8	2	20.0	2	18.2	7	30.4	4	12.9	3	7.1	15	22.1	3	11.5	6	25.0	5	21.7	3	21.4	1	14.3	
	200-220mg/dl未満	38	20.5	1	10.0	2	18.2	4	17.4	6	19.4	10	23.8	15	22.1	5	19.2	4	16.7	6	26.1	3	21.4	2	28.6	
	220-240mg/dl未満	34	18.4	1	10.0	1	9.1	2	8.7	8	25.8	12	28.6	10	14.7	8	30.8	4	16.7	4	17.4	2	14.3	-	-	
	240-260mg/dl未満	25	13.5	-	-	-	-	1	4.3	4	12.9	6	14.3	14	20.6	3	11.5	8	33.3	3	13.0	3	21.4	-	-	
	260-280mg/dl未満	14	7.6	-	-	-	-	1	4.3	2	6.5	7	16.7	4	5.9	4	15.4	1	4.2	1	4.3	2	14.3	-	-	
	280-300mg/dl未満	8	4.3	1	10.0	1	9.1	1	4.3	1	3.2	2	4.8	2	2.9	1	3.8	-	-	2	8.7	-	-	-	-	
	300mg/dl以上	2	1.1	-	-	-	-	-	-	1	3.2	1	2.4	-	-	1	3.8	-	-	-	-	-	-	-	-	

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬の服用者及び服用不明者除外

第 25 表の 4 総コレステロール値の平均値、標準偏差（20 歳以上、服薬者除外、性・年齢階級別）

(mg/dl)

年齢(歳)	男性			女性		
	人数	平均値	標準偏差	人数	平均値	標準偏差
総数	111	195.3	38.2	185	215.5	37.4
20-29歳	5	182.0	33.3	10	191.9	40.7
30-39歳	9	204.7	24.4	11	189.5	39.2
40-49歳	13	197.4	33.0	23	196.0	37.6
50-59歳	17	196.1	40.5	31	219.1	37.2
60-69歳	31	212.5	40.6	42	233.2	31.5
70歳以上	36	178.8	32.2	68	217.2	32.7
(再掲)						
65-69歳	19	219.0	43.7	26	231.0	34.2
70-74歳	10	187.4	14.8	24	222.4	27.0
75-79歳	13	176.8	35.7	23	218.8	31.9
80-84歳	7	182.1	37.9	14	225.9	31.3
85歳以上	6	164.7	32.6	7	176.6	26.9

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬の服用者及び服用不明者除外

第26表の1 HDL-コレステロール値の分布（20歳以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)										
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	65-69歳		70-74歳		75-79歳		80-84歳		85歳以上		
															人数	%	人数	%	人数	%	人数	%	人数	%	人数
総数	132	100.0	5	100.0	9	100.0	13	100.0	21	100.0	35	100.0	49	100.0	23	100.0	13	100.0	17	100.0	13	100.0	6	100.0	
男性	30mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	30-40mg/dl未満	21	15.9	1	20.0	-	1	7.7	4	19.0	2	5.7	13	26.5	2	8.7	3	23.1	5	29.4	3	23.1	2	33.3	
	40-50mg/dl未満	26	19.7	1	20.0	3	33.3	2	15.4	2	9.5	8	22.9	10	20.4	5	21.7	4	30.8	3	17.6	2	15.4	1	16.7
	50-60mg/dl未満	42	31.8	2	40.0	3	33.3	5	38.5	6	28.6	13	37.1	13	26.5	8	34.8	1	7.7	8	47.1	4	30.8	-	-
	60-70mg/dl未満	22	16.7	-	-	1	11.1	2	15.4	3	14.3	9	25.7	7	14.3	6	26.1	3	23.1	1	5.9	1	7.7	2	33.3
	70-80mg/dl未満	10	7.6	1	20.0	2	22.2	2	15.4	2	9.5	1	2.9	2	4.1	-	-	1	7.7	-	-	1	7.7	-	-
	80-90mg/dl未満	8	6.1	-	-	-	-	-	-	3	14.3	1	2.9	4	8.2	1	4.3	1	7.7	-	-	2	15.4	1	16.7
	90-100mg/dl未満	1	0.8	-	-	-	-	-	-	-	-	1	2.9	-	-	1	4.3	-	-	-	-	-	-	-	-
	100mg/dl以上	2	1.5	-	-	-	-	1	7.7	1	4.8	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	総数	248	100.0	10	100.0	12	100.0	23	100.0	38	100.0	66	100.0	99	100.0	38	100.0	31	100.0	36	100.0	20	100.0	12	100.0
女性	30mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	30-40mg/dl未満	7	2.8	-	-	-	-	-	-	-	2	3.0	5	5.1	1	2.6	-	-	1	2.8	2	10.0	2	16.7	
	40-50mg/dl未満	22	8.9	1	10.0	1	8.3	1	4.3	3	7.9	8	12.1	8	8.1	6	15.8	3	9.7	3	8.3	1	5.0	1	8.3
	50-60mg/dl未満	64	25.8	2	20.0	-	-	3	13.0	7	18.4	15	22.7	37	37.4	8	21.1	10	32.3	13	36.1	9	45.0	5	41.7
	60-70mg/dl未満	69	27.8	5	50.0	5	41.7	5	21.7	12	31.6	18	27.3	24	24.2	13	34.2	6	19.4	12	33.3	4	20.0	2	16.7
	70-80mg/dl未満	45	18.1	-	-	4	33.3	6	26.1	7	18.4	16	24.2	12	12.1	8	21.1	5	16.1	5	13.9	1	5.0	1	8.3
	80-90mg/dl未満	24	9.7	1	10.0	-	-	7	30.4	5	13.2	4	6.1	7	7.1	1	2.6	5	16.1	1	2.8	1	5.0	-	-
	90-100mg/dl未満	9	3.6	-	-	1	8.3	1	4.3	2	5.3	1	1.5	4	4.0	-	-	2	6.5	-	-	1	5.0	1	8.3
	100mg/dl以上	8	3.2	1	10.0	1	8.3	-	-	2	5.3	2	3.0	2	2.0	1	2.6	-	-	1	2.8	1	5.0	-	-

第26表の2 HDL-コレステロール値の平均値、標準偏差（20歳以上、性・年齢階級別）

(mg/dl)

年齢(歳)	男性			女性		
	人数	平均値	標準偏差	人数	平均値	標準偏差
総数	132	55.4	14.8	248	65.8	15.6
20-29歳	5	53.0	12.7	10	67.5	15.7
30-39歳	9	57.7	10.3	12	73.4	14.6
40-49歳	13	58.0	16.4	23	71.5	11.7
50-59歳	21	59.9	18.9	38	69.2	14.8
60-69歳	35	56.6	12.0	66	65.0	15.3
70歳以上	49	51.7	14.2	99	62.5	15.9
(再掲)						
65-69歳	23	56.6	13.1	38	62.9	15.7
70-74歳	13	53.6	15.1	31	65.6	14.1
75-79歳	17	48.1	8.0	36	63.1	16.1
80-84歳	13	54.1	16.5	20	60.0	17.1
85歳以上	6	53.0	18.1	12	56.8	15.4

第 26 表の 3 HDL-コレステロール値の分布（20 歳以上、服薬者除外、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)										
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	65-69歳		70-74歳		75-79歳		80-84歳		85歳以上		
															人数	%	人数	%	人数	%	人数	%	人数	%	人数
総数	111	100.0	5	100.0	9	100.0	13	100.0	17	100.0	31	100.0	36	100.0	19	100.0	10	100.0	13	100.0	7	100.0	6	100.0	
男性	30mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	30-40mg/dl未満	19	17.1	1	20.0	-	1	7.7	4	23.5	2	6.5	11	30.6	2	10.5	2	20.0	4	30.8	3	42.9	2	33.3	
	40-50mg/dl未満	22	19.8	1	20.0	3	33.3	2	15.4	-	-	8	25.8	8	22.2	5	26.3	4	40.0	2	15.4	1	14.3	1	16.7
	50-60mg/dl未満	35	31.5	2	40.0	3	33.3	5	38.5	5	29.4	12	38.7	8	22.2	7	36.8	-	-	6	46.2	2	28.6	-	-
	60-70mg/dl未満	20	18.0	-	-	1	11.1	2	15.4	3	17.6	7	22.6	7	19.4	4	21.1	3	30.0	1	7.7	1	14.3	2	33.3
	70-80mg/dl未満	8	7.2	1	20.0	2	22.2	2	15.4	1	5.9	1	3.2	1	2.8	-	-	1	10.0	-	-	-	-	-	
	80-90mg/dl未満	5	4.5	-	-	-	-	-	-	3	17.6	1	3.2	1	2.8	1	5.3	-	-	-	-	-	-	1	16.7
	90-100mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	100mg/dl以上	2	1.8	-	-	-	-	1	7.7	1	5.9	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	総数	185	100.0	10	100.0	11	100.0	23	100.0	31	100.0	42	100.0	68	100.0	26	100.0	24	100.0	23	100.0	14	100.0	7	100.0
女性	30mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	30-40mg/dl未満	5	2.7	-	-	-	-	-	-	-	1	2.4	4	5.9	1	3.8	-	-	1	4.3	2	14.3	1	14.3	
	40-50mg/dl未満	19	10.3	1	10.0	1	9.1	1	4.3	3	9.7	6	14.3	7	10.3	4	15.4	3	12.5	3	13.0	-	-	1	14.3
	50-60mg/dl未満	46	24.9	2	20.0	-	-	3	13.0	5	16.1	11	26.2	25	36.8	6	23.1	8	33.3	7	30.4	6	42.9	4	57.1
	60-70mg/dl未満	51	27.6	5	50.0	5	45.5	5	21.7	10	32.3	10	23.8	16	23.5	8	30.8	3	12.5	9	39.1	3	21.4	1	14.3
	70-80mg/dl未満	29	15.7	-	-	3	27.3	6	26.1	4	12.9	11	26.2	5	7.4	6	23.1	4	16.7	1	4.3	-	-	-	-
	80-90mg/dl未満	21	11.4	1	10.0	-	-	7	30.4	5	16.1	2	4.8	6	8.8	1	3.8	4	16.7	1	4.3	1	7.1	-	-
	90-100mg/dl未満	8	4.3	-	-	1	9.1	1	4.3	2	6.5	1	2.4	3	4.4	-	-	2	8.3	-	-	1	7.1	-	-
	100mg/dl以上	6	3.2	1	10.0	1	9.1	-	-	2	6.5	-	-	2	2.9	-	-	-	-	1	4.3	1	7.1	-	-

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬の服用者と服用不明者除外

第 26 表の 4 HDL-コレステロール値の平均値、標準偏差（20 歳以上、服薬者除外、性・年齢階級別）

(mg/dl)

年齢(歳)	男性			女性		
	人数	平均値	標準偏差	人数	平均値	標準偏差
総数	111	54.7	14.3	185	65.8	15.8
20-29歳	5	53.0	12.7	10	67.5	15.7
30-39歳	9	57.7	10.3	11	72.9	15.1
40-49歳	13	58.0	16.4	23	71.5	11.7
50-59歳	17	61.4	19.5	31	70.3	15.8
60-69歳	31	55.0	10.7	42	62.8	12.6
70歳以上	36	49.6	12.6	68	62.4	17.4
(再掲)						
65-69歳	19	53.8	11.2	26	61.0	12.4
70-74歳	10	52.5	13.4	24	65.4	15.0
75-79歳	13	48.2	8.4	23	63.3	19.3
80-84歳	7	45.4	10.1	14	61.2	19.3
85歳以上	6	53.0	18.1	7	51.4	7.0

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬の服用者と服用不明者除外

第27表の1 LDL-コレステロール値（直接法）の分布（20歳以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)									
															65-69歳		70-74歳		75-79歳		80-84歳		85歳以上	
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	132	100.0	5	100.0	9	100.0	13	100.0	21	100.0	35	100.0	49	100.0	23	100.0	13	100.0	17	100.0	13	100.0	6	100.0
40mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-50mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-60mg/dl未満	3	2.3	-	-	-	-	-	-	-	-	-	-	3	6.1	-	-	-	-	2	11.8	1	7.7	-	-
60-70mg/dl未満	6	4.5	1	20.0	-	-	-	-	2	9.5	1	2.9	2	4.1	1	4.3	-	-	-	-	1	7.7	1	16.7
70-80mg/dl未満	6	4.5	-	-	-	-	1	7.7	-	-	1	2.9	4	8.2	1	4.3	1	7.7	1	5.9	-	-	2	33.3
80-90mg/dl未満	13	9.8	-	-	1	11.1	-	-	5	23.8	1	2.9	6	12.2	-	-	1	7.7	3	17.6	1	7.7	1	16.7
90-100mg/dl未満	17	12.9	1	20.0	1	11.1	2	15.4	4	19.0	2	5.7	7	14.3	1	4.3	2	15.4	3	17.6	1	7.7	1	16.7
100-110mg/dl未満	10	7.6	1	20.0	-	-	3	23.1	1	4.8	1	2.9	4	8.2	-	-	2	15.4	-	-	2	15.4	-	-
110-120mg/dl未満	17	12.9	1	20.0	1	11.1	-	-	2	9.5	7	20.0	6	12.2	4	17.4	4	30.8	1	5.9	1	7.7	-	-
120-130mg/dl未満	20	15.2	-	-	-	-	2	15.4	3	14.3	6	17.1	9	18.4	5	21.7	-	-	6	35.3	3	23.1	-	-
130-140mg/dl未満	10	7.6	-	-	-	-	-	-	1	4.8	5	14.3	4	8.2	3	13.0	2	15.4	-	-	1	7.7	1	16.7
140-150mg/dl未満	9	6.8	-	-	3	33.3	3	23.1	1	4.8	1	2.9	1	2.0	1	4.3	-	-	1	5.9	-	-	-	-
150-160mg/dl未満	5	3.8	-	-	3	33.3	-	-	-	-	2	5.7	-	-	2	8.7	-	-	-	-	-	-	-	-
160-170mg/dl未満	5	3.8	-	-	-	-	-	-	2	9.5	3	8.6	-	-	1	4.3	-	-	-	-	-	-	-	-
170-180mg/dl未満	5	3.8	1	20.0	-	-	1	7.7	-	-	2	5.7	1	2.0	2	8.7	-	-	-	-	1	7.7	-	-
180-190mg/dl未満	2	1.5	-	-	-	-	1	7.7	-	-	1	2.9	-	-	-	-	-	-	-	-	-	-	-	-
190-200mg/dl未満	1	0.8	-	-	-	-	-	-	-	-	-	-	1	2.0	-	-	-	-	-	-	1	7.7	-	-
200-210mg/dl未満	2	1.5	-	-	-	-	-	-	-	-	-	1	2.9	1	2.0	1	4.3	1	7.7	-	-	-	-	-
210-220mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
220mg/dl以上	1	0.8	-	-	-	-	-	-	-	-	1	2.9	-	-	1	4.3	-	-	-	-	-	-	-	-
総数	248	100.0	10	100.0	12	100.0	23	100.0	38	100.0	66	100.0	99	100.0	38	100.0	31	100.0	36	100.0	20	100.0	12	100.0
40mg/dl未満	1	0.4	-	-	-	-	-	-	1	2.6	-	-	-	-	-	-	-	-	-	-	-	-	-	-
40-50mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
50-60mg/dl未満	2	0.8	-	-	1	8.3	-	-	-	-	1	1.5	-	-	-	-	-	-	-	-	-	-	-	-
60-70mg/dl未満	3	1.2	-	-	-	-	1	4.3	-	-	-	-	2	2.0	-	-	-	-	-	-	1	5.0	1	8.3
70-80mg/dl未満	11	4.4	-	-	2	16.7	1	4.3	3	7.9	3	4.5	2	2.0	2	5.3	1	3.2	1	2.8	-	-	-	-
80-90mg/dl未満	20	8.1	-	-	1	8.3	5	21.7	1	2.6	5	7.6	8	8.1	3	7.9	2	6.5	2	5.6	2	10.0	2	16.7
90-100mg/dl未満	24	9.7	3	30.0	1	8.3	4	17.4	3	7.9	2	3.0	11	11.1	1	2.6	2	6.5	6	16.7	1	5.0	2	16.7
100-110mg/dl未満	21	8.5	1	10.0	3	25.0	2	8.7	1	2.6	4	6.1	10	10.1	2	5.3	3	9.7	5	13.9	1	5.0	1	8.3
110-120mg/dl未満	35	14.1	2	20.0	2	16.7	2	8.7	6	15.8	6	9.1	17	17.2	5	13.2	4	12.9	6	16.7	5	25.0	2	16.7
120-130mg/dl未満	32	12.9	2	20.0	1	8.3	3	13.0	6	15.8	8	12.1	12	12.1	6	15.8	5	16.1	4	11.1	2	10.0	1	8.3
130-140mg/dl未満	23	9.3	-	-	-	-	1	4.3	5	13.2	5	7.6	12	12.1	1	2.6	5	16.1	4	11.1	2	10.0	1	8.3
140-150mg/dl未満	27	10.9	1	10.0	-	-	2	8.7	2	5.3	13	19.7	9	9.1	7	18.4	3	9.7	3	8.3	2	10.0	1	8.3
150-160mg/dl未満	16	6.5	-	-	1	8.3	-	-	4	10.5	8	12.1	3	3.0	4	10.5	2	6.5	1	2.8	-	-	-	-
160-170mg/dl未満	10	4.0	-	-	-	-	1	4.3	2	5.3	1	1.5	6	6.1	1	2.6	4	12.9	2	5.6	-	-	-	-
170-180mg/dl未満	9	3.6	1	10.0	-	-	-	-	2	5.3	3	4.5	3	3.0	1	2.6	-	-	-	-	3	15.0	-	-
180-190mg/dl未満	3	1.2	-	-	-	-	-	-	1	2.6	1	1.5	1	1.0	1	2.6	-	-	-	-	1	5.0	-	-
190-200mg/dl未満	5	2.0	-	-	-	-	-	-	-	-	3	4.5	2	2.0	2	5.3	-	-	1	2.8	-	-	1	8.3
200-210mg/dl未満	3	1.2	-	-	-	-	-	-	-	-	2	3.0	1	1.0	1	2.6	-	-	1	2.8	-	-	-	-
210-220mg/dl未満	2	0.8	-	-	-	-	-	-	1	2.6	1	1.5	-	-	1	2.6	-	-	-	-	-	-	-	-
220mg/dl以上	1	0.4	-	-	-	-	1	4.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

第27表の2 LDL-コレステロール値（直接法）の平均値、標準偏差（20歳以上、性・年齢階級別）

(mg/dl)

年齢(歳)	男性			女性		
	人数	平均値	標準偏差	人数	平均値	標準偏差
総数	132	117.6	33.2	248	124.0	33.0
20-29歳	5	110.4	36.4	10	117.5	25.1
30-39歳	9	132.2	25.2	12	101.0	25.7
40-49歳	13	124.5	30.7	23	111.5	34.7
50-59歳	21	108.2	28.3	38	127.0	34.1
60-69歳	35	132.7	33.4	66	133.9	34.9
70歳以上	49	106.9	30.9	99	122.6	29.6
(再掲)						
65-69歳	23	135.0	35.4	38	134.4	35.5
70-74歳	13	114.9	30.2	31	124.7	25.0
75-79歳	17	100.9	25.2	36	121.6	28.9
80-84歳	13	114.5	36.1	20	126.8	33.8
85歳以上	6	90.3	23.7	12	113.3	33.1

第 27 表の 3 LDL-コレステロール値（直接法）の分布（20 歳以上、服薬者除外、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)											
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	65-69歳		70-74歳		75-79歳		80-84歳		85歳以上			
															人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	111	100.0	5	100.0	9	100.0	13	100.0	17	100.0	31	100.0	36	100.0	19	100.0	10	100.0	13	100.0	7	100.0	6	100.0		
男性	40mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	40-50mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
	50-60mg/dl未満	3	2.7	-	-	-	-	-	-	-	-	-	3	8.3	-	-	-	-	2	15.4	1	14.3	-	-		
	60-70mg/dl未満	4	3.6	1	20.0	-	-	-	-	2	11.8	-	-	1	2.8	-	-	-	-	-	-	-	1	16.7		
	70-80mg/dl未満	6	5.4	-	-	-	-	1	7.7	-	-	1	3.2	4	11.1	1	5.3	1	10.0	1	7.7	-	-	2	33.3	
	80-90mg/dl未満	11	9.9	-	-	1	11.1	-	-	5	29.4	1	3.2	4	11.1	-	-	-	2	15.4	1	14.3	1	16.7		
	90-100mg/dl未満	13	11.7	1	20.0	1	11.1	2	15.4	4	23.5	2	6.5	3	8.3	1	5.3	1	10.0	1	7.7	-	-	1	16.7	
	100-110mg/dl未満	9	8.1	1	20.0	-	-	3	23.1	1	5.9	1	3.2	3	8.3	-	-	2	20.0	-	-	1	14.3	-	-	
	110-120mg/dl未満	15	13.5	1	20.0	1	11.1	-	-	1	5.9	6	19.4	6	16.7	3	15.8	4	40.0	1	7.7	1	14.3	-	-	
	120-130mg/dl未満	14	12.6	-	-	-	-	2	15.4	1	5.9	5	16.1	6	16.7	4	21.1	-	-	5	38.5	1	14.3	-	-	
	130-140mg/dl未満	10	9.0	-	-	-	-	-	-	1	5.9	5	16.1	4	11.1	3	15.8	2	20.0	-	-	-	1	14.3	1	16.7
	140-150mg/dl未満	8	7.2	-	-	3	33.3	3	23.1	1	5.9	-	-	1	2.8	-	-	-	-	1	7.7	-	-	-	-	
	150-160mg/dl未満	5	4.5	-	-	3	33.3	-	-	-	-	2	6.5	-	-	2	10.5	-	-	-	-	-	-	-	-	
	160-170mg/dl未満	4	3.6	-	-	-	-	-	-	1	5.9	3	9.7	-	-	1	5.3	-	-	-	-	-	-	-	-	
	170-180mg/dl未満	5	4.5	1	20.0	-	-	1	7.7	-	-	2	6.5	1	2.8	2	10.5	-	-	-	-	1	14.3	-	-	
	180-190mg/dl未満	2	1.8	-	-	-	-	1	7.7	-	-	1	3.2	-	-	-	-	-	-	-	-	-	-	-	-	
	190-200mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	200-210mg/dl未満	1	0.9	-	-	-	-	-	-	-	-	1	3.2	-	-	1	5.3	-	-	-	-	-	-	-	-	
	210-220mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
	220mg/dl以上	1	0.9	-	-	-	-	-	-	-	-	1	3.2	-	-	1	5.3	-	-	-	-	-	-	-	-	
	総数	185	100.0	10	100.0	11	100.0	23	100.0	31	100.0	42	100.0	68	100.0	26	100.0	24	100.0	23	100.0	14	100.0	7	100.0	
	女性	40mg/dl未満	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
40-50mg/dl未満		-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
50-60mg/dl未満		1	0.5	-	-	1	9.1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		
60-70mg/dl未満		2	1.1	-	-	-	-	1	4.3	-	-	-	-	1	1.5	-	-	-	-	-	-	-	1	14.3		
70-80mg/dl未満		5	2.7	-	-	1	9.1	1	4.3	2	6.5	-	-	1	1.5	-	-	1	4.2	-	-	-	-	-		
80-90mg/dl未満		11	5.9	-	-	1	9.1	5	21.7	1	3.2	2	4.8	2	2.9	2	7.7	-	-	-	-	-	2	28.6		
90-100mg/dl未満		17	9.2	3	30.0	1	9.1	4	17.4	3	9.7	-	-	6	8.8	-	-	2	8.3	3	13.0	-	-	1	14.3	
100-110mg/dl未満		11	5.9	1	10.0	3	27.3	2	8.7	1	3.2	-	-	4	5.9	-	-	1	4.2	3	13.0	-	-	-		
110-120mg/dl未満		25	13.5	2	20.0	2	18.2	2	8.7	5	16.1	3	7.1	11	16.2	3	11.5	3	12.5	3	13.0	4	28.6	1	14.3	
120-130mg/dl未満		25	13.5	2	20.0	1	9.1	3	13.0	3	9.7	6	14.3	10	14.7	5	19.2	4	16.7	4	17.4	2	14.3	-	-	
130-140mg/dl未満		19	10.3	-	-	-	-	1	4.3	4	12.9	4	9.5	10	14.7	1	3.8	4	16.7	3	13.0	2	14.3	1	14.3	
140-150mg/dl未満		23	12.4	1	10.0	-	-	2	8.7	2	6.5	10	23.8	8	11.8	5	19.2	3	12.5	2	8.7	2	14.3	1	14.3	
150-160mg/dl未満		14	7.6	-	-	1	9.1	-	-	4	12.9	6	14.3	3	4.4	3	11.5	2	8.3	1	4.3	-	-	-	-	
160-170mg/dl未満		10	5.4	-	-	-	-	1	4.3	2	6.5	1	2.4	6	8.8	1	3.8	4	16.7	2	8.7	-	-	-	-	
170-180mg/dl未満		9	4.9	1	10.0	-	-	-	-	2	6.5	3	7.1	3	4.4	1	3.8	-	-	-	-	3	21.4	-	-	
180-190mg/dl未満		3	1.6	-	-	-	-	-	-	1	3.2	1	2.4	1	1.5	1	3.8	-	-	-	-	1	7.1	-	-	
190-200mg/dl未満		4	2.2	-	-	-	-	-	-	-	-	3	7.1	1	1.5	2	7.7	-	-	1	4.3	-	-	-	-	
200-210mg/dl未満		3	1.6	-	-	-	-	-	-	-	-	2	4.8	1	1.5	1	3.8	-	-	1	4.3	-	-	-	-	
210-220mg/dl未満		2	1.1	-	-	-	-	-	-	1	3.2	1	2.4	-	-	1	3.8	-	-	-	-	-	-	-	-	
220mg/dl以上		1	0.5	-	-	-	-	1	4.3	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-		

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬の服用者と服用不明者除外

第 27 表の 4 LDL-コレステロール値（直接法）の平均値、標準偏差（20 歳以上、服薬者除外、性・年齢階級別）

年齢(歳)	男性			女性		
	人数	平均値	標準偏差	人数	平均値	標準偏差
総数	111	117.7	33.0	185	130.1	32.7
20-29歳	5	110.4	36.4	10	117.5	25.1
30-39歳	9	132.2	25.2	11	103.3	25.6
40-49歳	13	124.5	30.7	23	111.5	34.7
50-59歳	17	102.2	26.8	31	132.1	32.4
60-69歳	31	135.5	33.2	42	148.6	29.5
70歳以上	36	104.6	27.3	68	130.2	28.6
(再掲)						
65-69歳	19	140.1	35.0	26	145.3	33.3
70-74歳	10	110.9	17.2	24	130.0	24.3
75-79歳	13	101.8	27.7	23	131.2	29.5
80-84歳	7	113.3	34.3	14	142.1	26.2
85歳以上	6	90.3	23.7	7	103.6	26.5

注) コレステロールを下げる薬または中性脂肪（トリグリセライド）を下げる薬の服用者と服用不明者除外

第28表 メタボリックシンドローム（内臓脂肪症候群）の状況（20歳以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)					
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	40-74歳		65-74歳		75歳以上	
															人数	%	人数	%	人数	%
総数	372	100.0	15	100.0	20	100.0	36	100.0	59	100.0	99	100.0	143	100.0	238	100.0	105	100.0	99	100.0
メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者	81	21.8	1	6.7	-	-	1	2.8	10	16.9	26	26.3	43	30.1	48	20.2	29	27.6	32	32.3
メタボリックシンドローム(内臓脂肪症候群)の予備群と考えられる者	45	12.1	1	6.7	1	5.0	4	11.1	7	11.9	15	15.2	17	11.9	31	13.0	14	13.3	12	12.1
上記以外	246	66.1	13	86.7	19	95.0	31	86.1	42	71.2	58	58.6	83	58.0	159	66.8	62	59.0	55	55.6
総数	130	100.0	5	100.0	9	100.0	13	100.0	21	100.0	35	100.0	47	100.0	82	100.0	36	100.0	34	100.0
メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者	43	33.1	1	20.0	-	-	1	7.7	4	19.0	13	37.1	24	51.1	22	26.8	15	41.7	20	58.8
メタボリックシンドローム(内臓脂肪症候群)の予備群と考えられる者	27	20.8	-	-	1	11.1	4	30.8	3	14.3	8	22.9	11	23.4	18	22.0	7	19.4	8	23.5
上記以外	60	46.2	4	80.0	8	88.9	8	61.5	14	66.7	14	40.0	12	25.5	42	51.2	14	38.9	6	17.6
総数	242	100.0	10	100.0	11	100.0	23	100.0	38	100.0	64	100.0	96	100.0	156	100.0	69	100.0	65	100.0
メタボリックシンドローム(内臓脂肪症候群)が強く疑われる者	38	15.7	-	-	-	-	-	-	6	15.8	13	20.3	19	19.8	26	16.7	14	20.3	12	18.5
メタボリックシンドローム(内臓脂肪症候群)の予備群と考えられる者	18	7.4	1	10.0	-	-	-	-	4	10.5	7	10.9	6	6.3	13	8.3	7	10.1	4	6.2
上記以外	186	76.9	9	90.0	11	(100.0)	23	100.0	28	73.7	44	68.8	71	74.0	117	75.0	48	69.6	49	75.4

(注) 妊婦除外、項目の基準はp.6参照
 腹囲計測・血圧測定・血液検査未実施者、薬服用不明者除外。

第 29 表の 1 腹囲区分別、血中脂質、血圧、血糖のいずれかのリスクを有する割合
(20 歳以上、性・年齢階級別)

		総数		リスク無し		リスク数:1		リスク数:2		リスク数:3		
		人数	%	人数	%	人数	%	人数	%	人数	%	
総 数	総数	372	100.0	87	23.4	145	39.0	96	25.8	44	11.8	
	85cm未満(男)/90cm未満(女)	237	100.0	78	32.9	100	42.2	43	18.1	16	6.8	
	85cm以上(男)/90cm以上(女)	135	100.0	9	6.7	45	33.3	53	39.3	28	20.7	
	20-29歳	12	100.0	10	83.3	2	16.7	-	-	-	-	
	85cm以上(男)/90cm以上(女)	3	100.0	1	33.3	1	33.3	1	33.3	-	-	
	30-39歳	18	100.0	18	100.0	-	-	-	-	-	-	
	85cm以上(男)/90cm以上(女)	2	100.0	1	50.0	1	50.0	-	-	-	-	
	40-49歳	28	100.0	18	64.3	9	32.1	1	3.6	-	-	
	85cm以上(男)/90cm以上(女)	8	100.0	3	37.5	4	50.0	1	12.5	-	-	
	50-59歳	41	100.0	12	29.3	22	53.7	6	14.6	1	2.4	
	85cm以上(男)/90cm以上(女)	18	100.0	1	5.6	7	38.9	8	44.4	2	11.1	
	60-69歳	55	100.0	9	16.4	26	47.3	14	25.5	6	10.9	
	85cm以上(男)/90cm以上(女)	44	100.0	3	6.8	15	34.1	19	43.2	7	15.9	
	70歳以上	83	100.0	11	13.3	41	49.4	22	26.5	9	10.8	
	85cm以上(男)/90cm以上(女)	60	100.0	-	-	17	28.3	24	40.0	19	31.7	
	(再掲)											
	40-74歳	85cm未満(男)/90cm未満(女)	152	100.0	43	28.3	74	48.7	27	17.8	8	5.3
	85cm以上(男)/90cm以上(女)	86	100.0	7	8.1	31	36.0	33	38.4	15	17.4	
	65-74歳	85cm未満(男)/90cm未満(女)	60	100.0	10	16.7	31	51.7	15	25.0	4	6.7
	85cm以上(男)/90cm以上(女)	45	100.0	2	4.4	14	31.1	19	42.2	10	22.2	
75歳以上	85cm未満(男)/90cm未満(女)	55	100.0	7	12.7	24	43.6	16	29.1	8	14.5	
85cm以上(男)/90cm以上(女)	44	100.0	-	-	12	27.3	19	43.2	13	29.5		
男 性	総数	130	100.0	24	18.5	50	38.5	39	30.0	17	13.1	
	85cm未満	53	100.0	17	32.1	23	43.4	11	20.8	2	3.8	
	85cm以上	77	100.0	7	9.1	27	35.1	28	36.4	15	19.5	
	20-29歳	3	100.0	1	33.3	2	66.7	-	-	-	-	
	85cm以上	2	100.0	1	50.0	-	-	1	50.0	-	-	
	30-39歳	7	100.0	7	100.0	-	-	-	-	-	-	
	85cm以上	2	100.0	1	50.0	1	50.0	-	-	-	-	
	40-49歳	6	100.0	4	66.7	2	33.3	-	-	-	-	
	85cm以上	7	100.0	2	28.6	4	57.1	1	14.3	-	-	
	50-59歳	14	100.0	2	14.3	9	64.3	2	14.3	1	7.1	
	85cm以上	7	100.0	-	-	3	42.9	3	42.9	1	14.3	
	60-69歳	11	100.0	1	9.1	5	45.5	5	45.5	-	-	
	85cm以上	24	100.0	3	12.5	8	33.3	11	45.8	2	8.3	
	70歳以上	12	100.0	2	16.7	5	41.7	4	33.3	1	8.3	
	85cm以上	35	100.0	-	-	11	31.4	12	34.3	12	34.3	
	(再掲)											
	40-74歳	85cm未満	37	100.0	9	24.3	18	48.6	9	24.3	1	2.7
	85cm以上	45	100.0	5	11.1	18	40.0	16	35.6	6	13.3	
	65-74歳	85cm未満	12	100.0	2	16.7	4	33.3	6	50.0	-	-
	85cm以上	24	100.0	2	8.3	7	29.2	10	41.7	5	20.8	
75歳以上	85cm未満	6	100.0	-	-	3	50.0	2	33.3	1	16.7	
85cm以上	28	100.0	-	-	8	28.6	11	39.3	9	32.1		
女 性	総数	242	100.0	63	26.0	95	39.3	57	23.6	27	11.2	
	90cm未満	184	100.0	61	33.2	77	41.8	32	17.4	14	7.6	
	90cm以上	58	100.0	2	3.4	18	31.0	25	43.1	13	22.4	
	20-29歳	9	100.0	9	100.0	-	-	-	-	-	-	
	90cm以上	1	100.0	-	-	1	100.0	-	-	-	-	
	30-39歳	11	100.0	11	100.0	-	-	-	-	-	-	
	90cm以上	-	100.0	-	-	-	-	-	-	-	-	
	40-49歳	22	100.0	14	63.6	7	31.8	1	4.5	-	-	
	90cm以上	1	100.0	1	100.0	-	-	-	-	-	-	
	50-59歳	27	100.0	10	37.0	13	48.1	4	14.8	-	-	
	90cm以上	11	100.0	1	9.1	4	36.4	5	45.5	1	9.1	
	60-69歳	44	100.0	8	18.2	21	47.7	9	20.5	6	13.6	
	90cm以上	20	100.0	-	-	7	35.0	8	40.0	5	25.0	
	70歳以上	71	100.0	9	12.7	36	50.7	18	25.4	8	11.3	
	90cm以上	25	100.0	-	-	6	24.0	12	48.0	7	28.0	
	(再掲)											
	40-74歳	90cm未満	115	100.0	34	29.6	56	48.7	18	15.7	7	6.1
	90cm以上	41	100.0	2	4.9	13	31.7	17	41.5	9	22.0	
	65-74歳	90cm未満	48	100.0	8	16.7	27	56.3	9	18.8	4	8.3
	90cm以上	21	100.0	-	-	7	33.3	9	42.9	5	23.8	
75歳以上	90cm未満	49	100.0	7	14.3	21	42.9	14	28.6	7	14.3	
90cm以上	16	100.0	-	-	4	25.0	8	50.0	4	25.0		

第29表の2 腹囲区分別、血中脂質、血圧、血糖のリスクの保有状況
 [すべての組み合わせ①血中脂質②血圧③血糖] (20歳以上、性・年齢階級別)

		総数		①有②有③有		①有②有③無		①有②無③有		①有②無③無		①無②有③有		①無②有③無		①無②無③有		①無②無③無		
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	
総数	総数	372	100.0	44	11.8	38	10.2	5	1.3	10	2.7	53	14.2	122	32.8	13	3.5	87	23.4	
		85cm未満(男)/90cm未満(女)	237	100.0	16	6.8	16	6.8	3	1.3	9	3.8	24	10.1	81	34.2	10	4.2	78	32.9
		85cm以上(男)/90cm以上(女)	135	100.0	28	20.7	22	16.3	2	1.5	1	0.7	29	21.5	41	30.4	3	2.2	9	6.7
	20-29歳	85cm未満(男)/90cm未満(女)	12	100.0	-	-	-	-	-	-	-	-	-	-	2	16.7	-	-	10	83.3
		85cm以上(男)/90cm以上(女)	3	100.0	-	-	1	33.3	-	-	-	-	-	-	1	33.3	-	-	1	33.3
	30-39歳	85cm未満(男)/90cm未満(女)	18	100.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	18	100.0
		85cm以上(男)/90cm以上(女)	2	100.0	-	-	-	-	-	-	-	-	-	-	1	50.0	-	-	1	50.0
	40-49歳	85cm未満(男)/90cm未満(女)	28	100.0	-	-	-	-	-	-	-	-	1	3.6	6	21.4	3	10.7	18	64.3
		85cm以上(男)/90cm以上(女)	8	100.0	-	-	1	12.5	-	-	-	-	-	-	4	50.0	-	-	3	37.5
	50-59歳	85cm未満(男)/90cm未満(女)	41	100.0	1	2.4	3	7.3	-	-	2	4.9	3	7.3	18	43.9	2	4.9	12	29.3
		85cm以上(男)/90cm以上(女)	18	100.0	2	11.1	3	16.7	2	11.1	1	5.6	3	16.7	5	27.8	1	5.6	1	5.6
	60-69歳	85cm未満(男)/90cm未満(女)	55	100.0	6	10.9	6	10.9	-	-	3	5.5	8	14.5	22	40.0	1	1.8	9	16.4
		85cm以上(男)/90cm以上(女)	44	100.0	7	15.9	6	13.6	-	-	-	-	13	29.5	15	34.1	-	-	3	6.8
	70歳以上	85cm未満(男)/90cm未満(女)	83	100.0	9	10.8	7	8.4	3	3.6	4	4.8	12	14.5	33	39.8	4	4.8	11	13.3
		85cm以上(男)/90cm以上(女)	60	100.0	19	31.7	11	18.3	-	-	-	-	13	21.7	15	25.0	2	3.3	-	-
	(再掲)	85cm未満(男)/90cm未満(女)	152	100.0	8	5.3	11	7.2	-	-	7	4.6	16	10.5	60	39.5	7	4.6	43	28.3
		85cm以上(男)/90cm以上(女)	86	100.0	15	17.4	11	12.8	2	2.3	1	1.2	20	23.3	28	32.6	2	2.3	7	8.1
	65-74歳	85cm未満(男)/90cm未満(女)	60	100.0	4	6.7	6	10.0	-	-	4	6.7	9	15.0	26	43.3	1	1.7	10	16.7
		85cm以上(男)/90cm以上(女)	45	100.0	10	22.2	5	11.1	-	-	-	-	14	31.1	13	28.9	1	2.2	2	4.4
75歳以上	85cm未満(男)/90cm未満(女)	55	100.0	8	14.5	5	9.1	3	5.5	2	3.6	8	14.5	19	34.5	3	5.5	7	12.7	
	85cm以上(男)/90cm以上(女)	44	100.0	13	29.5	10	22.7	-	-	-	-	9	20.5	11	25.0	1	2.3	-	-	
男性	総数	130	100.0	17	13.1	16	12.3	1	0.8	2	1.5	22	16.9	46	35.4	2	1.5	24	18.5	
		85cm未満	53	100.0	2	3.8	3	5.7	-	-	1	1.9	8	15.1	21	39.6	1	1.9	17	32.1
		85cm以上	77	100.0	15	19.5	13	16.9	1	1.3	1	1.3	14	18.2	25	32.5	1	1.3	7	9.1
	20-29歳	85cm未満	3	100.0	-	-	-	-	-	-	-	-	-	-	2	66.7	-	-	1	33.3
		85cm以上	2	100.0	-	-	1	50.0	-	-	-	-	-	-	-	-	-	-	1	50.0
	30-39歳	85cm未満	7	100.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	7	100.0
		85cm以上	2	100.0	-	-	-	-	-	-	-	-	-	-	1	50.0	-	-	1	50.0
	40-49歳	85cm未満	6	100.0	-	-	-	-	-	-	-	-	-	-	2	33.3	-	-	4	66.7
		85cm以上	7	100.0	-	-	1	14.3	-	-	-	-	-	-	4	57.1	-	-	2	28.6
	50-59歳	85cm未満	14	100.0	1	7.1	1	7.1	-	-	-	-	1	7.1	9	64.3	-	-	2	14.3
		85cm以上	7	100.0	1	14.3	2	28.6	1	14.3	1	14.3	-	-	2	28.6	-	-	-	-
	60-69歳	85cm未満	11	100.0	-	-	1	9.1	-	-	-	-	4	36.4	4	36.4	1	9.1	1	9.1
		85cm以上	24	100.0	2	8.3	3	12.5	-	-	-	-	8	33.3	8	33.3	-	-	3	12.5
	70歳以上	85cm未満	12	100.0	1	8.3	1	8.3	-	-	1	8.3	3	25.0	4	33.3	-	-	2	16.7
		85cm以上	35	100.0	12	34.3	6	17.1	-	-	-	-	6	17.1	10	28.6	1	2.9	-	-
	(再掲)	85cm未満	37	100.0	1	2.7	3	8.1	-	-	-	-	6	16.2	17	45.9	1	2.7	9	24.3
		85cm以上	45	100.0	6	13.3	7	15.6	1	2.2	1	2.2	8	17.8	16	35.6	1	2.2	5	11.1
	65-74歳	85cm未満	12	100.0	-	-	2	16.7	-	-	-	-	4	33.3	4	33.3	-	-	2	16.7
		85cm以上	24	100.0	5	20.8	4	16.7	-	-	-	-	6	25.0	6	25.0	1	4.2	2	8.3
75歳以上	85cm未満	6	100.0	1	16.7	-	-	-	-	1	16.7	2	33.3	2	33.3	-	-	-	-	
	85cm以上	28	100.0	9	32.1	5	17.9	-	-	-	-	6	21.4	8	28.6	-	-	-	-	
女性	総数	242	100.0	27	11.2	22	9.1	4	1.7	8	3.3	31	12.8	76	31.4	11	4.5	63	26.0	
		90cm未満	184	100.0	14	7.6	13	7.1	3	1.6	8	4.3	16	8.7	60	32.6	9	4.9	61	33.2
		90cm以上	58	100.0	13	22.4	9	15.5	1	1.7	-	-	15	25.9	16	27.6	2	3.4	2	3.4
	20-29歳	90cm未満	9	100.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	9	100.0
		90cm以上	1	100.0	-	-	-	-	-	-	-	-	-	-	1	100.0	-	-	-	-
	30-39歳	90cm未満	11	100.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	11	100.0
		90cm以上	-	100.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	40-49歳	90cm未満	22	100.0	-	-	-	-	-	-	-	-	1	4.5	4	18.2	3	13.6	14	63.6
		90cm以上	1	100.0	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	100.0
	50-59歳	90cm未満	27	100.0	-	-	2	7.4	-	-	2	7.4	2	7.4	9	33.3	2	7.4	10	37.0
		90cm以上	11	100.0	1	9.1	1	9.1	1	9.1	-	-	3	27.3	3	27.3	1	9.1	1	9.1
	60-69歳	90cm未満	44	100.0	6	13.6	5	11.4	-	-	3	6.8	4	9.1	18	40.9	-	-	8	18.2
		90cm以上	20	100.0	5	25.0	3	15.0	-	-	-	-	5	25.0	7	35.0	-	-	-	-
	70歳以上	90cm未満	71	100.0	8	11.3	6	8.5	3	4.2	3	4.2	9	12.7	29	40.8	4	5.6	9	12.7
		90cm以上	25	100.0	7	28.0	5	20.0	-	-	-	-	7	28.0	5	20.0	1	4.0	-	-
	(再掲)	90cm未満	115	100.0	7	6.1	8	7.0	-	-	7	6.1	10	8.7	43	37.4	6	5.2	34	29.6
		90cm以上	41	100.0	9	22.0	4	9.8	1	2.4	-	-	12	29.3	12	29.3	1	2.4	2	4.9
	65-74歳	90cm未満	48	100.0	4	8.3	4	8.3	-	-	4	8.3	5	10.4	22	45.8	1	2.1	8	16.7
		90cm以上	21	100.0	5	23.8	1	4.8	-	-	-	-	8	38.1	7	33.3	-	-	-	-
75歳以上	90cm未満	49	100.0	7	14.3	5	10.2	3	6.1	1	2.0	6	12.2	17	34.7	3	6.1	7	14.3	
	90cm以上	16	100.0	4	25.0	5	31.3	-	-	-	-	3	18.8	3	18.8	1	6.3	-	-	

第30表 「糖尿病が強く疑われる者」及び「糖尿病の可能性を否定できない者」の状況（20歳以上、性・年齢階級別）

	総数	20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)						
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	40-74歳		65-74歳		75歳以上		
														人数	%	人数	%	人数	%	人数
総数	374	100.0	15	100.0	20	100.0	36	100.0	59	100.0	101	100.0	143	100.0	240	100.0	105	100.0	99	100.0
糖尿病が強く疑われる者	49	13.1	-	-	-	-	-	-	2	3.4	14	13.9	33	23.1	28	11.7	21	20.0	21	21.2
糖尿病が強く疑われる者のうち服薬者※	18	36.7	-	-	-	-	-	-	-	-	6	42.9	12	36.4	9	32.1	7	33.3	9	42.9
糖尿病の可能性を否定できない者	68	18.2	-	-	-	4	11.1	12	20.3	22	21.8	30	21.0	45	18.8	21	20.0	23	23.2	
上記以外	257	68.7	15	100.0	20	100.0	32	88.9	45	76.3	65	64.4	80	55.9	167	69.6	63	60.0	55	55.6
総数	128	100.0	5	100.0	9	100.0	13	100.0	21	100.0	35	100.0	45	100.0	82	100.0	36	100.0	32	100.0
糖尿病が強く疑われる者	24	18.7	-	-	-	-	-	-	2	9.5	8	22.9	14	31.1	14	17.1	11	30.6	10	31.3
糖尿病が強く疑われる者のうち服薬者※	8	33.3	-	-	-	-	-	-	-	-	3	37.5	5	35.7	3	21.4	3	27.3	5	50.0
糖尿病の可能性を否定できない者	20	15.6	-	-	-	-	-	-	2	9.5	8	22.9	10	22.2	13	15.9	8	22.2	7	21.9
上記以外	84	65.6	5	100.0	9	100.0	13	100.0	17	81.0	19	54.3	21	46.7	55	67.1	17	47.2	15	46.9
総数	246	100.0	10	100.0	11	100.0	23	100.0	38	100.0	66	100.0	98	100.0	158	100.0	69	100.0	67	100.0
糖尿病が強く疑われる者	25	10.2	-	-	-	-	-	-	-	-	9	9.1	19	19.4	14	8.9	10	14.5	11	16.4
糖尿病が強く疑われる者のうち服薬者※	10	40.0	-	-	-	-	-	-	-	-	3	50.0	7	36.8	6	42.9	4	40.0	4	36.4
糖尿病の可能性を否定できない者	48	19.5	-	-	-	4	17.4	10	26.3	14	21.2	20	20.4	32	20.3	13	18.8	16	23.9	
上記以外	173	70.3	10	100.0	11	100.0	19	82.6	28	73.7	46	69.7	59	60.2	112	70.9	46	66.7	40	59.7

※ 「糖尿病が強く疑われる者」に対する割合

「糖尿病が強く疑われる者」、「糖尿病の可能性を否定できない者」の判定

- ① 「糖尿病が強く疑われる者」とは、ヘモグロビンA1cの値が6.5%以上、または、問診で「現在糖尿病の治療を受けている」と答えた人である。
 ② 「糖尿病の可能性を否定できない者」とは、ヘモグロビンA1cの値が6.0%以上、6.5%未満で、①以外の人である。

第31表 高血圧症有病者の状況（20歳以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)					
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	40-74歳		65-74歳		75歳以上	
															人数	%	人数	%	人数	%
総数	438	100.0	19	100.0	22	100.0	37	100.0	64	100.0	115	100.0	181	100.0	278	100.0	129	100.0	119	100.0
高血圧症有病者	239	54.6	1	5.3	1	4.5	5	13.5	27	42.2	76	66.1	129	71.3	148	53.2	82	63.6	89	74.8
高血圧症有病者のうち服薬者※	138	57.7	-	-	-	-	-	-	12	44.4	34	44.7	92	71.3	73	49.3	45	54.9	65	73.0
正常高値血圧者	70	16.0	3	15.8	2	9.1	8	21.6	13	20.3	22	19.1	22	12.2	54	19.4	26	20.2	11	9.2
上記以外	129	29.5	15	78.9	19	86.4	24	64.9	24	37.5	17	14.8	30	16.6	76	27.3	21	16.3	19	16.0
総数	159	100.0	6	100.0	10	100.0	13	100.0	22	100.0	44	100.0	64	100.0	99	100.0	47	100.0	44	100.0
高血圧症有病者	95	59.7	1	16.7	1	10.0	3	23.1	12	54.5	30	68.2	48	75.0	56	56.6	29	61.7	37	84.1
高血圧症有病者のうち服薬者※	59	62.1	-	-	-	-	-	-	7	58.3	16	53.3	36	75.0	31	55.4	19	65.5	28	75.7
正常高値血圧者	32	20.1	2	33.3	1	10.0	4	30.8	6	27.3	9	20.5	10	15.6	24	24.2	12	25.5	5	11.4
上記以外	32	20.1	3	50.0	8	80.0	6	46.2	4	18.2	5	11.4	6	9.4	19	19.2	6	12.8	2	4.5
総数	279	100.0	13	100.0	12	100.0	24	100.0	42	100.0	71	100.0	117	100.0	179	100.0	82	100.0	75	100.0
高血圧症有病者	144	51.6	-	-	-	-	2	8.3	15	35.7	46	64.8	81	69.2	92	51.4	53	64.6	52	69.3
高血圧症有病者のうち服薬者※	79	54.9	-	-	-	-	-	-	5	33.3	18	39.1	56	69.1	42	45.7	26	49.1	37	71.2
正常高値血圧者	38	13.6	1	7.7	1	8.3	4	16.7	7	16.7	13	18.3	12	10.3	30	16.8	14	17.1	6	8.0
上記以外	97	34.8	12	92.3	11	91.7	18	75.0	20	47.6	12	16.9	24	20.5	57	31.8	15	18.3	17	22.7

※「高血圧症有病者」に対する割合

「高血圧症有病者」の判定

・高血圧症有病者とは、収縮期血圧が140mmHg以上、または、拡張期血圧が90mmHg以上、もしくは問診で「血圧を下げる薬」を使用していると答えた人である。

第32表 「脂質異常症が疑われる人」の状況（20歳以上、性・年齢階級別）

	総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)					
	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	40-74歳		65-74歳		75歳以上			
													人数	%	人数	%	人数	%	人数	%
総数	377	100.0	15	100.0	20	100.0	36	100.0	59	100.0	100	100.0	147	100.0	239	100.0	105	100.0	103	100.0
脂質異常症が疑われる人	105	27.9	1	6.7	-	-	1	2.8	15	25.4	30	30.0	58	39.5	58	24.3	31	29.5	46	44.7
脂質異常症が疑われる人のうち服薬者※	81	77.1	-	-	-	-	-	-	11	73.3	27	90.0	43	74.1	48	82.8	26	83.9	33	71.7
上記以外	272	72.1	14	93.3	20	100.0	35	97.2	44	74.6	70	70.0	89	60.5	181	75.7	74	70.5	57	55.3
総数	131	100.0	5	100.0	9	100.0	13	100.0	21	100.0	35	100.0	48	100.0	82	100.0	36	100.0	35	100.0
脂質異常症が疑われる人	39	29.8	1	20.0	-	-	1	7.7	8	38.1	6	17.1	23	47.9	20	24.4	11	30.6	18	51.4
脂質異常症が疑われる人のうち服薬者※	20	51.3	-	-	-	-	-	-	4	50.0	4	66.7	12	52.2	11	55.0	7	63.6	9	50.0
上記以外	92	70.2	4	80.0	9	100.0	12	92.3	13	61.9	29	82.9	25	52.1	62	75.6	25	69.4	17	48.6
総数	246	100.0	10	100.0	11	100.0	23	100.0	38	100.0	65	100.0	99	100.0	157	100.0	69	100.0	68	100.0
脂質異常症が疑われる人	66	26.8	-	-	-	-	-	-	7	18.4	24	36.9	35	35.4	38	24.2	20	29.0	28	41.2
脂質異常症が疑われる人のうち服薬者※	61	92.4	-	-	-	-	-	-	7	100.0	23	95.8	31	88.6	37	97.4	19	95.0	24	85.7
上記以外	180	73.2	10	100.0	11	100.0	23	100.0	31	81.6	41	63.1	64	64.6	119	75.8	49	71.0	40	58.8

※ 「脂質異常症が疑われる人」に対する割合

「脂質異常症が疑われる人」の判定

HDLコレステロールが40mg/dL未満、もしくは問診でコレステロールを下げる薬または中性脂肪を下げる薬を服用していると答えた人である。

第33表 1週間の運動日数（20歳以上、性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総数	408	100.0	19	100.0	21	100.0	32	100.0	61	100.0	107	100.0	168	100.0
	運動なし	190	46.6	11	57.9	16	76.2	26	81.3	40	65.6	46	43.0	51	30.4
	1日/週	38	9.3	4	21.1	-	-	1	3.1	6	9.8	10	9.3	17	10.1
	2日/週	44	10.8	2	10.5	-	-	2	6.3	4	6.6	16	15.0	20	11.9
	3日/週	31	7.6	2	10.5	2	9.5	1	3.1	3	4.9	6	5.6	17	10.1
	4日/週	13	3.2	-	-	-	-	-	-	-	-	4	3.7	9	5.4
	5日/週	19	4.7	-	-	1	4.8	-	-	3	4.9	5	4.7	10	6.0
	6日/週	8	2.0	-	-	-	-	-	-	1	1.6	3	2.8	4	2.4
7日/週	65	15.9	-	-	2	9.5	2	6.3	4	6.6	17	15.9	40	23.8	
男性	総数	147	100.0	6	100.0	10	100.0	11	100.0	20	100.0	39	100.0	61	100.0
	運動なし	66	44.9	4	66.7	6	60.0	10	90.9	14	70.0	17	43.6	15	24.6
	1日/週	14	9.5	1	16.7	-	-	-	-	3	15.0	3	7.7	7	11.5
	2日/週	17	11.6	-	-	-	-	1	9.1	3	15.0	6	15.4	7	11.5
	3日/週	10	6.8	1	16.7	1	10.0	-	-	-	-	-	-	8	13.1
	4日/週	4	2.7	-	-	-	-	-	-	-	-	1	2.6	3	4.9
	5日/週	10	6.8	-	-	1	10.0	-	-	-	-	2	5.1	7	11.5
	6日/週	2	1.4	-	-	-	-	-	-	-	-	-	-	2	3.3
7日/週	24	16.3	-	-	2	20.0	-	-	-	-	10	25.6	12	19.7	
女性	総数	261	100.0	13	100.0	11	100.0	21	100.0	41	100.0	68	100.0	107	100.0
	運動なし	124	47.5	7	53.8	10	90.9	16	76.2	26	63.4	29	42.6	36	33.6
	1日/週	24	9.2	3	23.1	-	-	1	4.8	3	7.3	7	10.3	10	9.3
	2日/週	27	10.3	2	15.4	-	-	1	4.8	1	2.4	10	14.7	13	12.1
	3日/週	21	8.0	1	7.7	1	9.1	1	4.8	3	7.3	6	8.8	9	8.4
	4日/週	9	3.4	-	-	-	-	-	-	-	-	3	4.4	6	5.6
	5日/週	9	3.4	-	-	-	-	-	-	3	7.3	3	4.4	3	2.8
	6日/週	6	2.3	-	-	-	-	-	-	1	2.4	3	4.4	2	1.9
7日/週	41	15.7	-	-	-	-	2	9.5	4	9.8	7	10.3	28	26.2	

第34表 運動を行う日の平均運動時間（20歳以上、性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上	
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%
総数	総数	218	100.0	8	100.0	5	100.0	6	100.0	21	100.0	61	100.0	117	100.0
	30分未満	37	17.0	-	-	1	20.0	3	50.0	1	4.8	10	16.4	22	18.8
	30分以上60分未満	83	38.1	4	50.0	1	20.0	1	16.7	8	38.1	27	44.3	42	35.9
	60分以上120分未満	64	29.4	1	12.5	1	20.0	1	16.7	9	42.9	20	32.8	32	27.4
	120分以上180分未満	24	11.0	3	37.5	2	40.0	1	16.7	2	9.5	3	4.9	13	11.1
	180分以上	10	4.6	-	-	-	-	-	-	1	4.8	1	1.6	8	6.8
男性	総数	81	100.0	2	100.0	4	100.0	1	100.0	6	100.0	22	100.0	46	100.0
	30分未満	7	8.6	-	-	-	-	-	-	-	-	2	9.1	5	10.9
	30分以上60分未満	28	34.6	2	100.0	1	25.0	-	-	-	-	9	40.9	16	34.8
	60分以上120分未満	26	32.1	-	-	1	25.0	1	100.0	4	66.7	10	45.5	10	21.7
	120分以上180分未満	12	14.8	-	-	2	50.0	-	-	1	16.7	1	4.5	8	17.4
	180分以上	8	9.9	-	-	-	-	-	-	1	16.7	-	-	7	15.2
女性	総数	137	100.0	6	100.0	1	100.0	5	100.0	15	100.0	39	100.0	71	100.0
	30分未満	30	21.9	-	-	1	100.0	3	60.0	1	6.7	8	20.5	17	23.9
	30分以上60分未満	55	40.1	2	33.3	-	-	1	20.0	8	53.3	18	46.2	26	36.6
	60分以上120分未満	38	27.7	1	16.7	-	-	-	-	5	33.3	10	25.6	22	31.0
	120分以上180分未満	12	8.8	3	50.0	-	-	1	20.0	1	6.7	2	5.1	5	7.0
	180分以上	2	1.5	-	-	-	-	-	-	-	-	1	2.6	1	1.4

注) 週1回以上運動を行うと回答した者

第35表 運動習慣の有無（20歳以上、性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	65-74歳		75歳以上	
																人数	%	人数	%
総数	総数	408	100.0	19	100.0	21	100.0	32	100.0	61	100.0	107	100.0	168	100.0	121	100.0	110	100.0
	運動習慣あり	128	31.4	3	15.8	4	19.0	2	6.3	13	21.3	34	31.8	72	42.9	49	40.5	47	42.7
	習慣なし	280	68.6	16	84.2	17	81.0	30	93.8	48	78.7	73	68.2	96	57.1	72	59.5	63	57.3
男性	総数	147	100.0	6	100.0	10	100.0	11	100.0	20	100.0	39	100.0	61	100.0	45	100.0	41	100.0
	運動習慣あり	56	38.1	1	16.7	4	40.0	1	9.1	3	15.0	15	38.5	32	52.5	20	44.4	23	56.1
	習慣なし	91	61.9	5	83.3	6	60.0	10	90.9	17	85.0	24	61.5	29	47.5	25	55.6	18	43.9
女性	総数	261	100.0	13	100.0	11	100.0	21	100.0	41	100.0	68	100.0	107	100.0	76	100.0	69	100.0
	運動習慣あり	72	27.6	2	15.4	-	-	1	4.8	10	24.4	19	27.9	40	37.4	29	38.2	24	34.8
	習慣なし	189	72.4	11	84.6	11	100.0	20	95.2	31	75.6	49	72.1	67	62.6	47	61.8	45	65.2

第36表の1 歩数の分布（20歳以上、性・年齢階級別）

		総数		20-29歳		30-39歳		40-49歳		50-59歳		60-69歳		70歳以上		(再掲)			
		人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	人数	%	20-64歳		65歳以上	
																人数	%	人数	%
総数	総数	586	100.0	37	100.0	40	100.0	73	100.0	84	100.0	138	100.0	214	100.0	290	100.0	296	100.0
	2000歩未満	72	12.3	2	5.4	2	5.0	5	6.8	6	7.1	10	7.2	47	22.0	18	6.2	54	18.2
	2000~3999歩	135	23.0	7	18.9	6	15.0	8	11.0	15	17.9	37	26.8	62	29.0	49	16.9	86	29.1
	4000~5999歩	137	23.4	11	29.7	6	15.0	22	30.1	23	27.4	31	22.5	44	20.6	81	27.9	56	18.9
	6000~7999歩	107	18.3	7	18.9	8	20.0	16	21.9	19	22.6	24	17.4	33	15.4	57	19.7	50	16.9
	8000~9999歩	77	13.1	6	16.2	9	22.5	12	16.4	13	15.5	22	15.9	15	7.0	49	16.9	28	9.5
	10000歩以上	58	9.9	4	10.8	9	22.5	10	13.7	8	9.5	14	10.1	13	6.1	36	12.4	22	7.4
	総数	253	100.0	18	100.0	20	100.0	32	100.0	37	100.0	57	100.0	89	100.0	127	100.0	126	100.0
男性	2000歩未満	34	13.4	1	5.6	1	5.0	2	6.3	4	10.8	7	12.3	19	21.3	10	7.9	24	19.0
	2000~3999歩	52	20.6	4	22.2	3	15.0	3	9.4	7	18.9	13	22.8	22	24.7	21	16.5	31	24.6
	4000~5999歩	48	19.0	3	16.7	-	-	9	28.1	7	18.9	11	19.3	18	20.2	27	21.3	21	16.7
	6000~7999歩	46	18.2	3	16.7	5	25.0	3	9.4	8	21.6	11	19.3	16	18.0	22	17.3	24	19.0
	8000~9999歩	36	14.2	3	16.7	6	30.0	5	15.6	7	18.9	7	12.3	8	9.0	22	17.3	14	11.1
	10000歩以上	37	14.6	4	22.2	5	25.0	10	31.3	4	10.8	8	14.0	6	6.7	25	19.7	12	9.5
	総数	333	100.0	19	100.0	20	100.0	41	100.0	47	100.0	81	100.0	125	100.0	163	100.0	170	100.0
女性	2000歩未満	38	11.4	1	5.3	1	5.0	3	7.3	2	4.3	3	3.7	28	22.4	8	4.9	30	17.6
	2000~3999歩	83	24.9	3	15.8	3	15.0	5	12.2	8	17.0	24	29.6	40	32.0	28	17.2	55	32.4
	4000~5999歩	89	26.7	8	42.1	6	30.0	13	31.7	16	34.0	20	24.7	26	20.8	54	33.1	35	20.6
	6000~7999歩	61	18.3	4	21.1	3	15.0	13	31.7	11	23.4	13	16.0	17	13.6	35	21.5	26	15.3
	8000~9999歩	41	12.3	3	15.8	3	15.0	7	17.1	6	12.8	15	18.5	7	5.6	27	16.6	14	8.2
	10000歩以上	21	6.3	-	-	4	20.0	-	-	4	8.5	6	7.4	7	5.6	11	6.7	10	5.9

第36表の2 歩数の平均値、標準偏差（20歳以上、性・年齢階級別）

年齢(歳)	総数			男性			女性		
	人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差
総数	586	5,676	3,542	253	6,154	4,143	333	5,312	2,961
20-29歳	37	6,762	5,121	18	8,159	6,770	19	5,438	2,316
30-39歳	40	7,564	4,005	20	8,343	4,719	20	6,784	3,063
40-49歳	73	6,910	3,493	32	8,131	4,344	41	5,957	2,283
50-59歳	84	6,020	3,134	37	6,038	3,504	47	6,006	2,848
60-69歳	138	5,922	3,330	57	6,023	3,736	81	5,851	3,034
70歳以上	214	4,420	2,988	89	4,678	3,064	125	4,237	2,932
(再掲)									
20-64歳	290	6,551	3,690	127	7,185	4,533	163	6,056	2,783
65歳以上	296	4,819	3,169	126	5,115	3,424	170	4,599	2,958

第36表の3 運動習慣の有無別の歩数の平均値、標準偏差（20歳以上、性・年齢階級別）

年齢(歳)		総数			男性			女性		
		人数	平均値	標準偏差	人数	平均値	標準偏差	人数	平均値	標準偏差
習慣あり	総数	118	5,640	2,941	51	5,617	3,205	67	5,657	2,747
	20-29歳	2	5,214	386	1	5,600	-	1	4,828	-
	30-39歳	3	6,282	2,663	3	6,282	2,663	-	-	-
	40-49歳	2	7,165	3,835	1	11,000	-	1	3,330	-
	50-59歳	13	5,516	3,045	3	4,916	5,597	10	5,696	2,291
	60-69歳	32	6,519	2,660	14	6,020	2,734	18	6,907	2,611
	70歳以上	66	5,175	3,020	29	5,241	3,282	37	5,124	2,842
習慣なし	総数	254	5,631	3,283	78	6,625	3,840	176	5,190	2,943
	20-29歳	15	7,601	5,565	5	10,374	8,974	10	6,214	2,444
	30-39歳	15	7,633	3,268	6	8,245	2,988	9	7,226	3,555
	40-49歳	29	6,435	3,010	9	8,897	3,194	20	5,328	2,213
	50-59歳	47	6,031	3,228	17	6,334	3,678	30	5,860	2,997
	60-69歳	66	5,712	3,513	19	6,521	4,769	47	5,385	2,856
	70歳以上	82	4,326	3,174	22	4,717	3,378	60	4,182	3,113

